

533 Rec'd PCT/PTO 10 AUG 2001

Form PTO-1390
(REV 11-98)

U.S. DEPARTMENT OF COMMERCE
PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

7126-2

**TRANSMITTAL LETTER TO THE UNITED STATES
DESIGNATED/ELECTED OFFICE (DO/EO/US)
CONCERNING A FILING UNDER 35 U.S.C. 371**

US APPLICATION NO. (If known, see 37 CFR 1.51)
09/913401

INTERNATIONAL APPLICATION NO.
PCT/CA99/00700

INTERNATIONAL FILING DATE
3 August 1999

PRIORITY DATE CLAIMED
17 February 1999

TITLE OF INVENTION
IMMOBILIZED LABELING COMPOUNDS AND METHODS

APPLICANT(S) FOR DO/EO/US

Alfred POLLAK; David ROE; Linda Fung LU; John THORNBAC; Catherine Michelle POLLOCK

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☒ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US)
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☒ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)) (unsigned)
10. ☐ A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A **FIRST** preliminary amendment.
☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information: 2 return receipt postcards, Article 34 Amendment, Int. Pre. Exam. Report

EXPRESS MAIL LABEL NO. EL 740156328US

ATTORNEY'S DOCKET NUMBER
7126-2

P1019158:1

Additionally the following defects have been observed:

- Additional claim fees of **\$116** as a non-small entity, including any required multiple dependent claim fee, are required. Applicant must submit the additional claim fees or cancel the additional claims for which fees are due.

SUMMARY OF FEES DUE:

Total additional fees required for this application is **\$246** for a Large Entity:

- **\$130** Late oath or declaration Surcharge.
- Total additional claim fee(s) for this application is **\$116**
 - **\$80** for 1 independent claims over 3.
 - **\$36** for 26 total claims over 20.

Applicant is reminded that any communications to the United States Patent and Trademark Office must be mailed to the address given in the heading and include the U.S. application no. shown above (37 CFR 1.5)

CHARITTA A BURT

Telephone: (703) 305-3734

PART 1 - ATTORNEY/APPLICANT COPY

U.S. APPLICATION NUMBER NO.	INTERNATIONAL APPLICATION NO	ATTY. DOCKET NO.
09/913,401	PCT/CA99/00700	7126-2

01/24/2002 MNGUYEN 00000125 09913401

01 FC:154	130.00 OP
02 FC:966	36.00 OP
03 FC:964	80.00 OP

Rec'd PCT/PTO 22 AUG 2001

09/913401

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:
Alfred Pollak
David Roe
Linda Fung Lu
John Thornback
Catherine M. Pollock

Serial No.: 09/913,401

Examiner: Unknown

Filing Date: August 10, 2001

Art Unit: Unknown

Atty. Dkt. No.: 7126-2

For: IMMOBILIZED LABELING COMPOUNDS AND METHODS

EXPRESS MAIL MAILING LABEL
NUMBER EL 740155319 US
DATE OF DEPOSIT August 22, 2001
I hereby certify that this paper or fee is being
deposited with the United State Postal Service
with sufficient postage "EXPRESS MAIL
POST
OFFICE TO ADDRESSEE" service under 37
C.F.R. 1.10 on the date indicated above and is
addressed to: Commissioner for
Patents, Washington D.C. 20231
Deanna Dolan
Deanna Dolan

PRELIMINARY AMENDMENT

Commissioner for Patents
Washington, DC 20231

Sir:

With respect to the above referenced patent application, Applicants respectfully request entry of this preliminary amendment.

If the check is inadvertently omitted, or should any additional fees under 37 C.F.R. §§ 1.16 to 1.21 be required for any reason relating to the enclosed materials, or should any overpayment be included herein, the Commissioner is authorized to deduct or credit said fees from or to Akerman, Senterfitt & Eidson, P.A. Deposit Account No. 50-0951.

AMENDMENT

In the Claims

Please amend claims 1-3, 7, 10-19, 21-22, 24, 26-29, 31-32, 35-40 and 42..

1. (Amended) A composition for generating a complex-forming metal ion labeled agent, the compositions comprising:
 - (a) a metal support surface; and
 - (b) a conjugate releasably bound to the support surface, the conjugate comprising a ligand and a targeting molecule;

wherein the conjugate [is capable of coordinating] coordinates with a complex-forming metal ion so that the labeled conjugate is released from the support surface.
2. (Amended) The composition of claim 1, wherein the metal support surface [is capable of] releasably [coordinating] coordinates to sulfur or phosphorous and the ligand comprises a sulfur or phosphorous atom for binding to the metal support surface.
3. (Amended) The composition of claim 2, wherein the ligand comprises a sulfur atom attached to a sulfur protecting group, wherein the metal support surface [being capable of binding] binds to the protected sulfur atom thereby releasing the sulfur protecting group from the sulfur atom and forming a thiol bond with the ligand.
7. (Amended) The composition of claim 2, wherein the ligand comprises:
 - (a) a surface binding group selected from the group consisting of a cysteine amino acid residue, a cysteine amino acid residue derivative, a thiol or thioester group attached to an organic molecule, an amino acid residue, an amino acid residue derivative including phosphorous and a phosphorous containing organic molecule, wherein the amino acid residue derivative or organic molecule [is capable of releasably binding] binds to the support surface; and
 - (b) at least one accessory group [capable of coordinating] coordinates with the complex-forming metal ion.
10. (Amended) The composition of [any one of claims 7 to 9] claim 8, wherein the ligand comprises 3 accessory groups selected from the group consisting of (a)

nitrogen, oxygen or sulfur atom incorporated in an amino acid residue; (b) nitrogen, oxygen, selenium, phosphorous or sulfur atom incorporated in an amino aside residue; (c) a nitrogen, oxygen, selenium, phosphorous or sulfur atom incorporated in an organic molecule; and (d) a combination of one or more of (a) to (c), wherein the residues, derivatives and/or molecules have metal coordinating activity.

11. (Amended) The composition of [any one of claims 1 to 10] claim 1, wherein the targeting molecule comprises a molecule having agonist or antagonist activity selected from the group consisting of a polypeptide, a peptide, a nucleic acid molecule, an oligonucleotide, a saccharide, an oligosaccharide, a steroid, a cyclic peptide, a peptide or polypeptide mimetic, an enzyme substrate, an inhibitor and a small organic molecule.
12. (Amended) The composition of [any one of claims 1 to 11] claim 1, wherein the targeting molecule comprises a peptide, a polypeptide, a peptide or polypeptide mimetic or a small organic molecule.
13. (Amended) The composition of [any one of claims 1 to 12] claim 1, wherein the targeting molecule comprises a molecule selected from the group consisting of a bombesin 7-14 fragment, QWAVGHLM, TKPPR, RGDS and a small organic molecule that targets a receptor or a transporter.
14. (Amended) The composition of [either of claims 6 or 13] claim 6 or claim 13, wherein the receptor or transporter is selected from the group consisting of a dopamine receptor or transporter, a serotonin receptor or transporter, a sigma receptor, GABA receptor, a nicotinic receptor, a cholinergic receptor, a norepinephrine receptor or transporter, a glucose transporter and an opiod receptor.
15. (Amended) The composition of [any one of claims 1 to 14] claim 3, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal [capable of releasably binding] that releasably binds sulfur or phosphorous for forming a metal complex.
16. (Amended) The composition of [any one of claims 1 to 14] claim 3, wherein the metal support surface comprises gold.
17. (Amended) The composition of [any one of claims 1 to 16] claim 15, wherein the complex-forming metal is selected from the group of metals and radioisotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.
18. (Amended) The composition of [any one of claims 1 to 16] claim 17, wherein the complex-forming metal is selected from the group of metals and radiosotopic metals consisting of Tc, Cu and Re.

- 209110-10421650
19. (Amended) A method for generating a complex-forming metal ion labeled diagnostic agent or radiotherapeutic agent, comprising: (a) providing a composition according to [any one of claims 1 to 18] claim 1; and (b) contacting the composition with the complex-forming metal ion to form a coordinate bond between the complex-forming metal ion and the agent so that the complex-forming metal labeled agent is released from the support surface.
 21. (Amended) A metal ion labeled agent prepared using a composition of [any one of claims 1 to 18] claim 1.
 22. (Amended) A technetium or rhenium labeled agent prepared using a composition of [any one of claims 1 to 18] claim 1, wherein the agent is labeled with ^{99m}Tc and has a specific activity of greater than 10,000 Ci/mmol or the agent is labeled with ^{188}Re and has a specific activity of greater than 3,000 Ci/mmol.
 24. (Amended) A pharmaceutical composition for radiotherapy or imaging, comprising a carrier and a complex-forming metal ion labeled agent, wherein the agent is prepared using a composition of [any one of claims 1 to 18] claim 1.
 26. (Amended) A method of detecting the presence or assessing the severity of a disease, disorder or abnormal physical state in a mammal comprising:
 - (a) administering an effective amount of the agent or composition of [any one of claims 21 to 25] claim 22 or claim 24; and
 - (b) detecting the presence or assessing the severity of the disease, disorder or abnormal physical state.
 27. (Amended) A method of radiotherapy of a disease, disorder or abnormal physical state in a mammal comprising administering an effective amount of the agent or composition of [any one of claims 21 to 25] claim 22 or claim 24.
 28. (Amended) The method of [either of claims 26 or 27] claim 26, wherein the complex-forming metal labeled imaging agent is administered by an intravenous route.
 29. (Amended) The method of [any one of claims 26 to 28] claim 26, wherein the amount of complex-forming metal labeled agent administered to the mammal is about 0.01 mcg/kg/minute to 1,000 mcg/kg/minute.
 31. (Amended) The method of [any one of claims 26 to 30] claim 26, wherein the mammal is a human.
 32. (Amended) The method of [any one of claims 26 to 31] claim 26, wherein the disease, disorder or abnormal physical state is selected from the group consisting of oncological, neurological, inflammatory, infection, and degenerative diseases, disorders and abnormal physical states.

- 209170-1041660
35. (Amended) A kit for preparing a complex-forming metal ion labeled agent, the kit comprising a metal support surface, conjugate and a predetermined quantity of complex-forming metal ion, the conjugate being [capable of being] releasably bound to the support surface and [capable of coordinating] which coordinates with the complex-forming metal ion so that the conjugate is released from the metal support surface.
 36. (Amended) The kit of claim 35, wherein the conjugate comprises a sulfur atom attached to a sulfur protecting group, wherein the metal support surface [being capable of binding] binds to the protected sulfur atom thereby releasing the sulfur protecting group from the sulfur atom and forming a thiol bond with the conjugate.
 37. (Amended) The kit of claim 35, wherein the metal support surface [is capable of] releasably [coordinating] coordinates to sulfur or phosphorous and the conjugate comprises a sulfur or phosphorous atom for binding to the metal support surface.
 38. (Amended) The kit of claim 27, wherein the conjugate comprises a ligand and a targeting molecule, wherein the ligand comprises:
 - (a) a surface binding group selected for the group consisting of a cysteine amino acid residue, a cysteine amino acid residue derivative, a thiol or thioester group attached to an organic molecule, an amino acid residue derivative including phosphorous and a phosphorous containing organic molecule, wherein the amino acid residue, amino acid residue derivative or organic molecule [is capable of releasably binding] releasably binds to the support surface; and
 - (b) at least one accessory group [capable of coordinating] that coordinates with the complex-forming metal ion.
 39. (Amended) The kit of [any one of claims 35 to 38] claim 35, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal [capable of releasably binding] that binds sulfur or phosphorous for forming a metal complex; and the complex-forming metal is selected from the group of metals and radiosotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.
 40. (Amended) The kit of [any one of claims 35 to 39] claim 39, further comprising at least one agent selected form the group consisting of a reducing agent, a bulking agent and a pH stabilising agent.
 42. (Amended) The method of claim 41, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal [capable of releasably binding] that releasably binds sulfur or [forming] forms a metal complex; and the complex-forming metal is selected from the group of metals and radioisotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.

REMARKS


Status of the Claims

Claims 1-3, 7, 10-19, 21-22, 24, 26-29, 31-32, 35-40 and 42 have been amended. Claims 1-44 are pending in the case.

A separate clean copy of the amended claims is attached as Appendix A.

Applicants have submitted this amendment in order to reduce the calculated number of dependent claims.

Respectfully submitted,



Barbara S. Kitchell
Reg. No. 33,928
AKERMAN SENTERFITT
222 Lakeview Avenue, Suite 400
P.O. Box 3188
West Palm Beach, Florida 33402-3188
Tel: 561.653.5123

APPENDIX A

1. A composition for generating a complex-forming metal ion labeled agent, the compositions comprising:
 - (a) a metal support surface; and
 - (b) a conjugate releasably bound to the support surface, the conjugate comprising a ligand and a targeting molecule;

wherein the conjugate coordinates with a complex-forming metal ion so that the labeled conjugate is released from the support surface.

2. The composition of claim 1, wherein the metal support surface releasably coordinates to sulfur or phosphorous and the ligand comprises a sulfur or phosphorous atom for binding to the metal support surface.
3. The composition of claim 2, wherein the ligand comprises a sulfur atom attached to a sulfur protecting group, wherein the metal support surface binds to the protected sulfur atom thereby releasing the sulfur protecting group from the sulfur atom and forming a thiol bond with the ligand.
4. The composition of claim 2, wherein the conjugate comprises a peptide, a polypeptide, a peptide or polypeptide mimetic or a small organic molecule.
5. The composition of claim 4, wherein the conjugate comprises a peptide sequence selected from the group consisting of a bombesin 7-14 fragment, QWAVGHLM, TKPPR and RGDS.
6. The composition of claim 4, wherein the conjugate comprises a small organic molecule that targets a receptor or a transporter.
7. The composition of claim 2, wherein the ligand comprises:
 - (a) a surface binding group selected from the group consisting of a cysteine amino acid residue, a cysteine amino acid residue derivative, a thiol or thioester group attached to an organic molecule, an amino acid residue, an amino acid residue derivative including phosphorous and a phosphorous containing organic molecule, wherein the amino acid residue derivative or organic molecule binds to the support surface; and
 - (b) at least one accessory group coordinates with the complex-forming metal ion.
8. The composition of claim 7, wherein the ligand comprises a peptide, a peptide mimetic, a polypeptide, a polypeptide mimetic or a small organic molecule.

9. The composition of claim 8, wherein the ligand comprises a peptide selected from the group consisting of a tetradentate N_xS_{4-x} ligand, a tetradentate N_xS_{4-x} ligand derivative, a polyamino polysulfide and a polyamino polysulfide derivative.
10. The composition of claim 8, wherein the ligand comprises 3 accessory groups selected from the group consisting of (a) nitrogen, oxygen or sulfur atom incorporated in an amino acid residue; (b) nitrogen, oxygen, selenium, phosphorous or sulfur atom incorporated in an amino aside residue; (c) a nitrogen, oxygen, selenium, phosphorous or sulfur atom incorporated in an organic molecule; and (d) a combination of one or more of (a) to (c), wherein the residues, derivatives and/or molecules have metal coordinating activity.
11. The composition of claim 1, wherein the targeting molecule comprises a molecule having agonist or antagonist activity selected from the group consisting of a polypeptide, a peptide, a nucleic acid molecule, an oligonucleotide, a saccharide, an oligosaccharide, a steroid, a cyclic peptide, a peptide or polypeptide mimetic, an enzyme substrate, an inhibitor and a small organic molecule.
12. The composition of claim 1, wherein the targeting molecule comprises a peptide, a polypeptide, a peptide or polypeptide mimetic or a small organic molecule.
13. The composition of claim 1, wherein the targeting molecule comprises a molecule selected from the group consisting of a bombesin 7-14 fragment, QWAVGHLM, TKPPR, RGDS and a small organic molecule that targets a receptor or a transporter.
14. The composition of claim 6 or claim 13, wherein the receptor or transporter is selected from the group consisting of a dopamine receptor or transporter, a serotonin receptor or transporter, a sigma receptor, GABA receptor, a nicotinic receptor, a cholinergic receptor, a norepinephrine receptor or transporter, a glucose transporter and an opiod receptor.
15. The composition of claim 3, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal that releasably binds sulfur or phosphorous for forming a metal complex.
16. The composition of, wherein the metal support surface comprises gold.
17. The composition of claim 15, wherein the complex-forming metal is selected from the group of metals and radioisotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.
18. The composition of claim 17, wherein the complex-forming metal is selected from the group of metals and radiosotopic metals consisting of Tc, Cu and Re.
19. A method for generating a complex-forming metal ion labeled diagnostic agent or radiotherapeutic agent, comprising: (a) providing a composition according to claim 1; and (b) contacting the composition with the complex-forming metal ion to form a

coordinate bond between the complex-forming metal ion and the agent so that the complex-forming metal labeled agent is released from the support surface.

20. The method of claim 19, further comprising collecting the complex-forming metal labeled agent so released.
21. A metal ion labeled agent prepared using a composition of claim 1.
22. A technetium or rhenium labeled agent prepared using a composition of claim 1, wherein the agent is labeled with ^{99m}Tc and has a specific activity of greater than 10,000 Ci/mmol or the agent is labeled with ^{188}Re and has a specific activity of greater than 3,000 Ci/mmol.
23. The composition of claim 22, wherein the agent is a peptide comprising dimethylglycylserinylcysteinylglycine.
24. A pharmaceutical composition for radiotherapy or imaging, comprising a carrier and a complex-forming metal ion labeled agent, wherein the agent is prepared using a composition of claim 1.
25. The pharmaceutical composition of claim 24 further comprising at least one agent selected from the group consisting of a reducing agent, a bulking agent and a pH stabilizing agent.
26. A method of detecting the presence or assessing the severity of a disease, disorder or abnormal physical state in a mammal comprising:
 - (a) administering an effective amount of the agent or composition of claim 22 or claim 24; and
 - (b) detecting the presence or assessing the severity of the disease, disorder or abnormal physical state.
27. A method of radiotherapy of a disease, disorder or abnormal physical state in a mammal comprising administering an effective amount of the agent or composition of claim 22 or claim 24.
28. The method of claim 26, wherein the complex-forming metal labeled imaging agent is administered by an intravenous route.
29. The method of claim 26, wherein the amount of complex-forming metal labeled agent administered to the mammal is about 0.01 mcg/kg/minute to 1,000 mcg/kg/minute.
30. The method of claim 29, wherein the amount of the complex-forming metal labeled agent administered to the mammal is about 0.01 to 50 mcg/kg/minute.
31. The method of claim 26, wherein the mammal is a human.

32. The method of claim 26, wherein the disease, disorder or abnormal physical state is selected from the group consisting of oncological, neurological, inflammatory, infection, and degenerative diseases, disorders and abnormal physical states.
33. The method of claim 32, wherein the presence or the severity of a disease, disorder or abnormal physical state is detected or assessed with a technique selected from the group consisting of positron emission tomography, nuclear magnetic resonance imaging, scintigraphy, single photon emission computed tomography, perfusion contrast echocardiography, ultrafast X-ray computed tomography, and digital subtraction angiography.
34. The method of claim 33, wherein the agent comprises a ^{99m}Tc metal and binds to a receptor and the technique is single photon emission computed tomography.
35. A kit for preparing a complex-forming metal ion labeled agent, the kit comprising a metal support surface, conjugate and a predetermined quantity of complex-forming metal ion, the conjugate being releasably bound to the support surface and which coordinates with the complex-forming metal ion so that the conjugate is released from the metal support surface.
36. The kit of claim 35, wherein the conjugate comprises a sulfur atom attached to a sulfur protecting group, wherein the metal support surface binds to the protected sulfur atom thereby releasing the sulfur protecting group from the sulfur atom and forming a thiol bond with the conjugate.
37. The kit of claim 35, wherein the metal support surface releasably coordinates to sulfur or phosphorous and the conjugate comprises a sulfur or phosphorous atom for binding to the metal support surface.
38. The kit of claim 27, wherein the conjugate comprises a ligand and a targeting molecule, wherein the ligand comprises:
- (a) a surface binding group selected for the group consisting of a cysteine amino acid residue, a cysteine amino acid residue derivative, a thiol or thioester group attached to an organic molecule, an amino acid residue derivative including phosphorous and a phosphorous containing organic molecule, wherein the amino acid residue, amino acid residue derivative or organic molecule releasably binds to the support surface; and
 - (b) at least one accessory group that coordinates with the complex-forming metal ion.
39. The kit of claim 35, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal that binds sulfur or phosphorous for forming a metal complex; and the complex-forming metal is selected from the group of metals and radiosotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.

- 209110-1041600
40. The kit of claim 39, further comprising at least one agent selected from the group consisting of a reducing agent, a bulking agent and a pH stabilising agent.
41. A method for generating a complex-forming metal ion labeled agent comprising:
- (a) providing a metal support surface;
 - (b) providing a conjugate comprising a ligand and targeting molecule, wherein the ligand comprises a peptide, a peptide mimetic, a polypeptide or a polypeptide mimetic of about 3 to 50 amino acid residues or derivatives thereof and includes a sulfur atom for binding to the metal support surface, the sulfur atom being protected by a sulfur protecting group;
 - (c) contacting the protected sulfur atom with the metal support surface so that the sulfur atom forms a thiol bond with the metal surface thereby releasing the sulfur protecting group; and
 - (d) contacting the ligand with the complex-forming metal ion to form a coordinate bond between the complex-forming metal ion and the ligand so that the complex-forming metal labeled agent is released from the support surface.
42. The method of claim 41, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal that releasably binds sulfur or forms a metal complex; and the complex-forming metal is selected from the group of metals and radioisotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.
43. A method for the preparation of a support surface for manufacturing a complex-forming metal labeled agent comprising electro or electroless metal plating or vapor deposition of a suitable thickness of the metal onto an inorganic or polymeric substrate in the form of particles, sponges or sieves, fibers or surfaces with suitable surface area between about 1 and 10,000 cm².
44. The method of claim 43, wherein the thickness of the metal on the support surface is greater than about 10nm.



08-23 - JC20 Rec'd PCT/PTO P2 AUG 2001
8650
5000
1097913401
Burt

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:
Alfred Pollak
David Roe
Linda Fung Lu
John Thornback
Catherine M. Pollock

Serial No.: 09/913,401

Examiner: Unknown

Filing Date: August 10, 2001

Art Unit: Unknown

Atty. Dkt. No.: 7126-2

For: IMMOBILIZED LABELING COMPOUNDS AND METHODS

EXPRESS MAIL MAILING LABEL
NUMBER EL 740155319 US
DATE OF DEPOSIT August 22, 2001
I hereby certify that this paper or fee is being
deposited with the United State Postal Service
with sufficient postage "EXPRESS MAIL
POST
OFFICE TO ADDRESSEE" service under 37
C.F.R. 1.10 on the date indicated above and is
addressed to: Commissioner for
Patents, Washington D.C. 20231

Deanna Dolan
Deanna Dolan

PRELIMINARY AMENDMENT

Commissioner for Patents
Washington, DC 20231

Sir:

With respect to the above referenced patent application, Applicants respectfully request
entry of this preliminary amendment.

If the check is inadvertently omitted, or should any additional fees under 37 C.F.R. §§ 1.16 to 1.21 be required for any reason relating to the enclosed materials, or should any overpayment be included herein, the Commissioner is authorized to deduct or credit said fees from or to Akerman, Senterfitt & Eidson, P.A. Deposit Account No. 50-0951.

AMENDMENT

In the Claims

Please amend claims 1-3, 7, 10-19, 21-22, 24, 26-29, 31-32, 35-40 and 42..

1. (Amended) A composition for generating a complex-forming metal ion labeled agent, the compositions comprising:
 - (a) a metal support surface; and
 - (b) a conjugate releasably bound to the support surface, the conjugate comprising a ligand and a targeting molecule;wherein the conjugate [is capable of coordinating] coordinates with a complex-forming metal ion so that the labeled conjugate is released from the support surface.
2. (Amended) The composition of claim 1, wherein the metal support surface [is capable of] releasably [coordinating] coordinates to sulfur or phosphorous and the ligand comprises a sulfur or phosphorous atom for binding to the metal support surface.
3. (Amended) The composition of claim 2, wherein the ligand comprises a sulfur atom attached to a sulfur protecting group, wherein the metal support surface [being capable of binding] binds to the protected sulfur atom thereby releasing the sulfur protecting group from the sulfur atom and forming a thiol bond with the ligand.
7. (Amended) The composition of claim 2, wherein the ligand comprises:
 - (a) a surface binding group selected from the group consisting of a cysteine amino acid residue, a cysteine amino acid residue derivative, a thiol or thioester group attached to an organic molecule, an amino acid residue, an amino acid residue derivative including phosphorous and a phosphorous containing organic molecule, wherein the amino acid residue derivative or organic molecule [is capable of releasably binding] binds to the support surface; and
 - (b) at least one accessory group [capable of coordinating] coordinates with the complex-forming metal ion.
10. (Amended) The composition of [any one of claims 7 to 9] claim 8, wherein the ligand comprises 3 accessory groups selected from the group consisting of (a)

nitrogen, oxygen or sulfur atom incorporated in an amino acid residue; (b) nitrogen, oxygen, selenium, phosphorous or sulfur atom incorporated in an amino aside residue; (c) a nitrogen, oxygen, selenium, phosphorous or sulfur atom incorporated in an organic molecule; and (d) a combination of one or more of (a) to (c), wherein the residues, derivatives and/or molecules have metal coordinating activity.

11. (Amended) The composition of [any one of claims 1 to 10] claim 1, wherein the targeting molecule comprises a molecule having agonist or antagonist activity selected from the group consisting of a polypeptide, a peptide, a nucleic acid molecule, an oligonucleotide, a saccharide, an oligosaccharide, a steroid, a cyclic peptide, a peptide or polypeptide mimetic, an enzyme substrate, an inhibitor and a small organic molecule.
12. (Amended) The composition of [any one of claims 1 to 11] claim 1, wherein the targeting molecule comprises a peptide, a polypeptide, a peptide or polypeptide mimetic or a small organic molecule.
13. (Amended) The composition of [any one of claims 1 to 12] claim 1, wherein the targeting molecule comprises a molecule selected from the group consisting of a bombesin 7-14 fragment, QWAVGHLM, TKPPR, RGDS and a small organic molecule that targets a receptor or a transporter.
14. (Amended) The composition of [either of claims 6 or 13] claim 6 or claim 13, wherein the receptor or transporter is selected from the group consisting of a dopamine receptor or transporter, a serotonin receptor or transporter, a sigma receptor, GABA receptor, a nicotinic receptor, a cholinergic receptor, a norepinephrine receptor or transporter, a glucose transporter and an opiod receptor.
15. (Amended) The composition of [any one of claims 1 to 14] claim 3, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal [capable of releasably binding] that releasably binds sulfur or phosphorous for forming a metal complex.
16. (Amended) The composition of [any one of claims 1 to 14] claim 3, wherein the metal support surface comprises gold.
17. (Amended) The composition of [any one of claims 1 to 16] claim 15, wherein the complex-forming metal is selected from the group of metals and radioisotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.
18. (Amended) The composition of [any one of claims 1 to 16] claim 17, wherein the complex-forming metal is selected from the group of metals and radiosotopic metals consisting of Tc, Cu and Re.

19. (Amended) A method for generating a complex-forming metal ion labeled diagnostic agent or radiotherapeutic agent, comprising: (a) providing a composition according to [any one of claims 1 to 18] claim 1; and (b) contacting the composition with the complex-forming metal ion to form a coordinate bond between the complex-forming metal ion and the agent so that the complex-forming metal labeled agent is released from the support surface.
21. (Amended) A metal ion labeled agent prepared using a composition of [any one of claims 1 to 18] claim 1.
22. (Amended) A technetium or rhenium labeled agent prepared using a composition of [any one of claims 1 to 18] claim 1, wherein the agent is labeled with ^{99m}Tc and has a specific activity of greater than 10,000 Ci/mmol or the agent is labeled with ^{188}Re and has a specific activity of greater than 3,000 Ci/mmol.
24. (Amended) A pharmaceutical composition for radiotherapy or imaging, comprising a carrier and a complex-forming metal ion labeled agent, wherein the agent is prepared using a composition of [any one of claims 1 to 18] claim 1.
26. (Amended) A method of detecting the presence or assessing the severity of a disease, disorder or abnormal physical state in a mammal comprising:
- (a) administering an effective amount of the agent or composition of [any one of claims 21 to 25] claim 22 or claim 24; and
 - (b) detecting the presence or assessing the severity of the disease, disorder or abnormal physical state.
27. (Amended) A method of radiotherapy of a disease, disorder or abnormal physical state in a mammal comprising administering an effective amount of the agent or composition of [any one of claims 21 to 25] claim 22 or claim 24.
28. (Amended) The method of [either of claims 26 or 27] claim 26, wherein the complex-forming metal labeled imaging agent is administered by an intravenous route.
29. (Amended) The method of [any one of claims 26 to 28] claim 26, wherein the amount of complex-forming metal labeled agent administered to the mammal is about 0.01 mcg/kg/minute to 1,000 mcg/kg/minute.
31. (Amended) The method of [any one of claims 26 to 30] claim 26, wherein the mammal is a human.
32. (Amended) The method of [any one of claims 26 to 31] claim 26, wherein the disease, disorder or abnormal physical state is selected from the group consisting of oncological, neurological, inflammatory, infection, and degenerative diseases, disorders and abnormal physical states.

- 209470-101660
35. (Amended) A kit for preparing a complex-forming metal ion labeled agent, the kit comprising a metal support surface, conjugate and a predetermined quantity of complex-forming metal ion, the conjugate being [capable of being] releasably bound to the support surface and [capable of coordinating] which coordinates with the complex-forming metal ion so that the conjugate is released from the metal support surface.
36. (Amended) The kit of claim 35, wherein the conjugate comprises a sulfur atom attached to a sulfur protecting group, wherein the metal support surface [being capable of binding] binds to the protected sulfur atom thereby releasing the sulfur protecting group from the sulfur atom and forming a thiol bond with the conjugate.
37. (Amended) The kit of claim 35, wherein the metal support surface [is capable of] releasably [coordinating] coordinates to sulfur or phosphorous and the conjugate comprises a sulfur or phosphorous atom for binding to the metal support surface.
38. (Amended) The kit of claim 27, wherein the conjugate comprises a ligand and a targeting molecule, wherein the ligand comprises:
- (a) a surface binding group selected for the group consisting of a cysteine amino acid residue, a cysteine amino acid residue derivative, a thiol or thioester group attached to an organic molecule, an amino acid residue derivative including phosphorous and a phosphorous containing organic molecule, wherein the amino acid residue, amino acid residue derivative or organic molecule [is capable of releasably binding] releasably binds to the support surface; and
 - (b) at least one accessory group [capable of coordinating] that coordinates with the complex-forming metal ion.
39. (Amended) The kit of [any one of claims 35 to 38] claim 35, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal [capable of releasably binding] that binds sulfur or phosphorous for forming a metal complex; and the complex-forming metal is selected from the group of metals and radiosotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.
40. (Amended) The kit of [any one of claims 35 to 39] claim 39, further comprising at least one agent selected from the group consisting of a reducing agent, a bulking agent and a pH stabilising agent.
42. (Amended) The method of claim 41, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal [capable of releasably binding] that releasably binds sulfur or [forming] forms a metal complex; and the complex-forming metal is selected from the group of metals and radioisotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.

REMARKS


Status of the Claims

Claims 1-3, 7, 10-19, 21-22, 24, 26-29, 31-32, 35-40 and 42 have been amended. Claims 1-44 are pending in the case.

A separate clean copy of the amended claims is attached as Appendix A.

Applicants have submitted this amendment in order to reduce the calculated number of dependent claims.

Respectfully submitted,



Barbara S. Kitchell

Reg. No. 33,928

AKERMAN SENTERFITT

222 Lakeview Avenue, Suite 400

P.O. Box 3188

West Palm Beach, Florida 33402-3188

Tel: 561.653.5123

APPENDIX A

1. A composition for generating a complex-forming metal ion labeled agent, the compositions comprising:
 - (a) a metal support surface; and
 - (b) a conjugate releasably bound to the support surface, the conjugate comprising a ligand and a targeting molecule;

wherein the conjugate coordinates with a complex-forming metal ion so that the labeled conjugate is released from the support surface.

2. The composition of claim 1, wherein the metal support surface releasably coordinates to sulfur or phosphorous and the ligand comprises a sulfur or phosphorous atom for binding to the metal support surface.
3. The composition of claim 2, wherein the ligand comprises a sulfur atom attached to a sulfur protecting group, wherein the metal support surface binds to the protected sulfur atom thereby releasing the sulfur protecting group from the sulfur atom and forming a thiol bond with the ligand.
4. The composition of claim 2, wherein the conjugate comprises a peptide, a polypeptide, a peptide or polypeptide mimetic or a small organic molecule.
5. The composition of claim 4, wherein the conjugate comprises a peptide sequence selected from the group consisting of a bombesin 7-14 fragment, QWAVGHLM, TKPPR and RGDS.
6. The composition of claim 4, wherein the conjugate comprises a small organic molecule that targets a receptor or a transporter.
7. The composition of claim 2, wherein the ligand comprises:
 - (a) a surface binding group selected from the group consisting of a cysteine amino acid residue, a cysteine amino acid residue derivative, a thiol or thioester group attached to an organic molecule, an amino acid residue, an amino acid residue derivative including phosphorous and a phosphorous containing organic molecule, wherein the amino acid residue derivative or organic molecule binds to the support surface; and
 - (b) at least one accessory group coordinates with the complex-forming metal ion.
8. The composition of claim 7, wherein the ligand comprises a peptide, a peptide mimetic, a polypeptide, a polypeptide mimetic or a small organic molecule.

9. The composition of claim 8, wherein the ligand comprises a peptide selected from the group consisting of a tetradentate N_xS_{4-x} ligand, a tetradentate N_xS_{4-x} ligand derivative, a polyamino polysulfide and a polyamino polysulfide derivative.
10. The composition of claim 8, wherein the ligand comprises 3 accessory groups selected from the group consisting of (a) nitrogen, oxygen or sulfur atom incorporated in an amino acid residue; (b) nitrogen, oxygen, selenium, phosphorous or sulfur atom incorporated in an amino aside residue; (c) a nitrogen, oxygen, selenium, phosphorous or sulfur atom incorporated in an organic molecule; and (d) a combination of one or more of (a) to (c), wherein the residues, derivatives and/or molecules have metal coordinating activity.
11. The composition of claim 1, wherein the targeting molecule comprises a molecule having agonist or antagonist activity selected from the group consisting of a polypeptide, a peptide, a nucleic acid molecule, an oligonucleotide, a saccharide, an oligosaccharide, a steroid, a cyclic peptide, a peptide or polypeptide mimetic, an enzyme substrate, an inhibitor and a small organic molecule.
12. The composition of claim 1, wherein the targeting molecule comprises a peptide, a polypeptide, a peptide or polypeptide mimetic or a small organic molecule.
13. The composition of claim 1, wherein the targeting molecule comprises a molecule selected from the group consisting of a bombesin 7-14 fragment, QWAVGHLM, TKPPR, RGDS and a small organic molecule that targets a receptor or a transporter.
14. The composition of claim 6 or claim 13, wherein the receptor or transporter is selected from the group consisting of a dopamine receptor or transporter, a serotonin receptor or transporter, a sigma receptor, GABA receptor, a nicotinic receptor, a cholinergic receptor, a norepinephrine receptor or transporter, a glucose transporter and an opiod receptor.
15. The composition of claim 3, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal that releasably binds sulfur or phosphorous for forming a metal complex.
16. The composition of, wherein the metal support surface comprises gold.
17. The composition of claim 15, wherein the complex-forming metal is selected from the group of metals and radioisotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.
18. The composition of claim 17, wherein the complex-forming metal is selected from the group of metals and radiosotopic metals consisting of Tc, Cu and Re.
19. A method for generating a complex-forming metal ion labeled diagnostic agent or radiotherapeutic agent, comprising: (a) providing a composition according to claim 1; and (b) contacting the composition with the complex-forming metal ion to form a

coordinate bond between the complex-forming metal ion and the agent so that the complex-forming metal labeled agent is released from the support surface.

20. The method of claim 19, further comprising collecting the complex-forming metal labeled agent so released.
21. A metal ion labeled agent prepared using a composition of claim 1.
22. A technetium or rhenium labeled agent prepared using a composition of claim 1, wherein the agent is labeled with ^{99m}Tc and has a specific activity of greater than 10,000 Ci/mmol or the agent is labeled with ^{188}Re and has a specific activity of greater than 3,000 Ci/mmol.
23. The composition of claim 22, wherein the agent is a peptide comprising dimethylglycylserinylcysteinylglycine.
24. A pharmaceutical composition for radiotherapy or imaging, comprising a carrier and a complex-forming metal ion labeled agent, wherein the agent is prepared using a composition of claim 1.
25. The pharmaceutical composition of claim 24 further comprising at least one agent selected from the group consisting of a reducing agent, a bulking agent and a pH stabilizing agent.
26. A method of detecting the presence or assessing the severity of a disease, disorder or abnormal physical state in a mammal comprising:
 - (a) administering an effective amount of the agent or composition of claim 22 or claim 24; and
 - (b) detecting the presence or assessing the severity of the disease, disorder or abnormal physical state.
27. A method of radiotherapy of a disease, disorder or abnormal physical state in a mammal comprising administering an effective amount of the agent or composition of claim 22 or claim 24.
28. The method of claim 26, wherein the complex-forming metal labeled imaging agent is administered by an intravenous route.
29. The method of claim 26, wherein the amount of complex-forming metal labeled agent administered to the mammal is about 0.01 mcg/kg/minute to 1,000 mcg/kg/minute.
30. The method of claim 29, wherein the amount of the complex-forming metal labeled agent administered to the mammal is about 0.01 to 50 mcg/kg/minute.
31. The method of claim 26, wherein the mammal is a human.

32. The method of claim 26, wherein the disease, disorder or abnormal physical state is selected from the group consisting of oncological, neurological, inflammatory, infection, and degenerative diseases, disorders and abnormal physical states.
33. The method of claim 32, wherein the presence or the severity of a disease, disorder or abnormal physical state is detected or assessed with a technique selected from the group consisting of positron emission tomography, nuclear magnetic resonance imaging, scintigraphy, single photon emission computed tomography, perfusion contrast echocardiography, ultrafast X-ray computed tomography, and digital subtraction angiography.
34. The method of claim 33, wherein the agent comprises a ^{99m}Tc metal and binds to a receptor and the technique is single photon emission computed tomography.
35. A kit for preparing a complex-forming metal ion labeled agent, the kit comprising a metal support surface, conjugate and a predetermined quantity of complex-forming metal ion, the conjugate being releasably bound to the support surface and which coordinates with the complex-forming metal ion so that the conjugate is released from the metal support surface.
36. The kit of claim 35, wherein the conjugate comprises a sulfur atom attached to a sulfur protecting group, wherein the metal support surface binds to the protected sulfur atom thereby releasing the sulfur protecting group from the sulfur atom and forming a thiol bond with the conjugate.
37. The kit of claim 35, wherein the metal support surface releasably coordinates to sulfur or phosphorous and the conjugate comprises a sulfur or phosphorous atom for binding to the metal support surface.
38. The kit of claim 27, wherein the conjugate comprises a ligand and a targeting molecule, wherein the ligand comprises:
- (a) a surface binding group selected for the group consisting of a cysteine amino acid residue, a cysteine amino acid residue derivative, a thiol or thioester group attached to an organic molecule, an amino acid residue derivative including phosphorous and a phosphorous containing organic molecule, wherein the amino acid residue, amino acid residue derivative or organic molecule releasably binds to the support surface; and
 - (b) at least one accessory group that coordinates with the complex-forming metal ion.
39. The kit of claim 35, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal that binds sulfur or phosphorous for forming a metal complex; and the complex-forming metal is selected from the group of metals and radiotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.

40. The kit of claim 39, further comprising at least one agent selected from the group consisting of a reducing agent, a bulking agent and a pH stabilising agent.
41. A method for generating a complex-forming metal ion labeled agent comprising:
- (a) providing a metal support surface;
 - (b) providing a conjugate comprising a ligand and targeting molecule, wherein the ligand comprises a peptide, a peptide mimetic, a polypeptide or a polypeptide mimetic of about 3 to 50 amino acid residues or derivatives thereof and includes a sulfur atom for binding to the metal support surface, the sulfur atom being protected by a sulfur protecting group;
 - (c) contacting the protected sulfur atom with the metal support surface so that the sulfur atom forms a thiol bond with the metal surface thereby releasing the sulfur protecting group; and
 - (d) contacting the ligand with the complex-forming metal ion to form a coordinate bond between the complex-forming metal ion and the ligand so that the complex-forming metal labeled agent is released from the support surface.
42. The method of claim 41, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal that releasably binds sulfur or forms a metal complex; and the complex-forming metal is selected from the group of metals and radioisotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.
43. A method for the preparation of a support surface for manufacturing a complex-forming metal labeled agent comprising electro or electroless metal plating or vapor deposition of a suitable thickness of the metal onto an inorganic or polymeric substrate in the form of particles, sponges or sieves, fibers or surfaces with suitable surface area between about 1 and 10,000 cm².
44. The method of claim 43, wherein the thickness of the metal on the support surface is greater than about 10nm.

IMMOBILIZED LABELING COMPOUNDS AND METHODSFIELD OF THE INVENTION

The invention relates to compounds and methods useful for generating metal-ligand complexes. The metal-labeled agents are useful in radiotherapy and as imaging agents in medical diagnostics.

BACKGROUND

The art of diagnostic imaging exploits compounds that in binding or localizing site selectively within the body, help to resolve the image of diagnostic interest. Diagnostic imaging agents are effective for targeting and imaging biological receptors. These diagnostic imaging agents incorporate radionuclide metals such as technetium and rhenium. The radionuclide metals are used to label targeting molecules, such as proteins, peptides and antibodies that localize at desired regions of the human body. Localization of these agents is detected by gamma camera analysis. As targeting agents, proteins and peptides can offer the tissue specificity required for diagnostic accuracy.

Labeling of these and other diagnostic agents with metal atoms is made difficult by their chemical structure. Conventional labeling techniques involve the formation of the metal complex in a solution of excess ligand which typically results in high levels of unlabelled ligand. For example, technetium labeling reactions yield approximately one labeled ligand for every thousand or more unlabeled ligand. For many radiodiagnostic agents there is a finite number of binding sites ie. receptors for which both the labeled and unlabeled diagnostic agent compete. This leads to poor imaging because the excess unlabeled ligands compete with the labeled ligands for the binding sites. As a result, the administration of larger doses of the agent is required to achieve an acceptable image. The administration of larger doses of the agent into a patient can lead to adverse effects. Targeting molecules often cause physiological changes when they bind or localize to specific sites. For example, the targeting molecule may be a peptide that binds to a receptor and agonizes or antagonizes receptor activity. Unlabeled peptide will still cause the physiological effect of agonizing or antagonizing receptor activity, however, the benefits of a metal labeled peptide will not be obtained.

Presently, high performance liquid chromatography (HPLC) is applied to enhance the concentration of labeled agent in the solution before it is administered. While enhancing concentration, this separation step requires additional time and expense. This makes the technique impractical for clinical use.

5

A method for labeling biological targeting molecules with radionuclide metals is described in US patent 5,789,555. The method involves the steps of covalently linking a chelating portion of a biological targeting molecule to a solid support through a metal cleavable maleimide linker. When a radionuclide metal is introduced, it complexes with the chelating
10 portion of the biological targeting molecule. This results in the release of the biological targeting molecule from the support yielding a labeled biological targeting molecule. This method provides a high ratio of labeled to unlabeled molecules (specific activity). However the use of the linker requires an extra step of coupling the linker to the support. The requirement of a linker is therefore a drawback of this method.

15

Many labeling methods require the formation of the free thiol. The thiols tend react with each other forming disulphides. Extra precautions need to be taken to avoid disulphide formation which results in the reduction of product attached to the surface and the need for extra purification before radiolabeling.

20

Many metal chelators contain a sulfur atom which participates in the chelating function. Sulfur atoms are commonly rendered unreactive by the use of protecting groups, one of which is acetamidomethyl. The sulfur is protected with an acetamidomethyl group. The
25 sulfur protecting groups must be removed to provide a free thiol. The acetamidomethyl protecting group is removed from the sulfur through the use of mercuric acetate. Mercuric acetate coordinates to the sulfur which then possesses a slightly positive charge making the chelator a good leaving group. The excess mercuric acetate is removed by reaction with hydrogen sulphide. Mercuric acetate is quite toxic. Therefore
30 the requirement to deprotect sulfur with mercuric acetate is a drawback.

The labeling method described in US patent 5,789,555 makes use of organic solid supports that consist of a plurality of particles. The particles can be difficult to filter for the subsequent
35 removal of the unbound and labeled biological targeting molecules. These supports are

very susceptible to radiolytic degradation due the emission of β -particles by certain radionuclide metals. These supports are also otherwise not chemically robust. This makes the solid support difficult to sterilize since sterilization is carried out at very high temperatures.

5

There is therefore a need for a solid support for labeling biological targeting molecules with a radionuclide metal that permits labeling be carried out in a single step without the need for linker group. There is a need for a solid support for labeling biological targeting molecules with a radionuclide metal that has increased chemical robustness, reduced radiolytic degradation, is easily sterilized, and labeled and has well defined loading characteristics. There is further need for such a solid support that can directly bind a protected thiol group to form a bound thiol thereby avoiding the formation of disulfides.

10

There is a further need for an improved system that employs a surface as a substitute for mercury so that no mercuric acetate needs to be used or subsequently removed in order to deprotect sulfur atoms. There is a need for such a system that eliminates a few synthetic steps as well as the presence of mercury in a reaction solution.

15

SUMMARY OF THE INVENTION

The invention includes a method of producing metal labeled imaging and radiopharmaceutical agent formulations having high specific activity. The invention includes labeling peptides (such as dimethylglycylserinylcysteinylglycine) or other organic molecules that are attached to a support surface, preferably gold, via a group of a molecule that binds the surface. The surface binding molecule is preferably the sulfur group of a cysteine (this cysteine will ultimately be a part of the metal chelate). During the labeling of the peptide with a metal, (which is preferably a radioisotope, such as ^{99m}Tc), the complexation of the metal to the chelator, and hence to the sulfur of the cysteine, causes weakening of the gold-cysteine bond with the result that the metal complexed peptide leaves the surface and moves into solution. Uncomplexed peptides remain attached to the gold surface. The result of this selective cleavage from the surface is that the radiolabeled peptide is produced without added carrier giving a high specific activity formulation.

20

25

30

According to one aspect of the present invention, sulfur is initially de-protected to produce the free thiol using mercuric acetate. The de-protected peptide is then added to the gold. However, the gold surface also allows for the *in situ* formation of the bound thiol. Therefore, according to another aspect of the present invention, a novel method for

5 *in situ* production of the bound thiol is employed. This method allows protected sulfur groups to be de-protected without having to use mercuric acetate.

This approach to the preparation of radiopharmaceuticals having high specific activity offers the following advantages over the use of polymer supported strategies:

- 10
- Increased chemical robustness of the metal support surface
 - Elimination of the presence of mercuric acetate
 - Avoidance of unnecessary handling of the free thiol
 - Elimination of extra synthetic steps
- 15
- Production of the bound thiol *in situ*
 - Reduced radiolytic degradation of the metal support surface
 - Ease of sterilisation
- 20
- Ease of attachment of the HS-chelator-targeting molecule
 - Well defined loading characteristics (by virtue of the ability to vary the surface area)

The invention includes a compound useful for generating a complex-forming metal ion labeled agent, the compound including: a metal support surface, and a conjugate releasably bound to the support surface, wherein the conjugate is capable of

25 coordinating with a complex-forming metal ion so that the conjugate is released from the support surface. In a preferred variant, the conjugate includes a ligand and a targeting molecule and the ligand incorporates:

5 (a) a surface binding group selected from the group consisting of a cysteine amino acid residue, a cysteine amino acid residue derivative, a thiol or thioether group attached to an organic molecule, an amino acid residue derivative including phosphorous and a phosphorous containing organic molecule, wherein the amino acid residue, amino acid residue derivative or organic molecule is capable of releasably binding to the support surface; and

10 (b) at least one accessory group capable of coordinating with the complex-forming metal ion; wherein the conjugate is capable of coordinating with a complex-forming metal ion so that the conjugate is released from the support surface.

15 In another embodiment, the invention includes a method for generating a complex-forming metal ion labeled diagnostic agent or radiotherapeutic agent, including the steps of: (a) obtaining a compound according to any of claims 1 to 18; (b) contacting the compound with the complex-forming metal ion to form a coordinate bond between the complex-forming metal ion and the agent so that the complex-forming metal labeled agent is released from the support surface.

20 One variation of the invention includes a method for generating a complex-forming metal ion labeled diagnostic agent or radiotherapeutic agent, including transchelating a conjugate from a metal support surface to a complex-forming metal ion so that the conjugate is released from the metal support surface. The invention also includes a method for preparing a complex-forming metal ion labeled peptide including (a) bonding a peptide, polypeptide, peptide mimetic or polypeptide mimetic that includes (i) a cysteine amino acid residue, a cysteine amino acid residue derivative or a phosphorous amino acid residue derivative and (ii) at least one accessory group capable of coordinating with the complex-forming metal ion, to a metal support surface by the sulfur atom or phosphorous atom of the residue or derivative and (b) labeling the peptide, polypeptide or mimetic with a complex-forming metal so that the metal complexed peptide, polypeptide or mimetic is released from the support surface. Another aspect of the invention relates to a method for preparing a complex-forming metal ion labeled agent including (a) bonding an organic molecule that includes (i) a sulfur atom or a phosphorous atom and (ii) at least one accessory group capable of coordinating with the complex-forming metal ion, to a support surface by the sulfur atom or the phosphorous atom and (b) labeling the agent with a

complex-forming metal so that the metal complex conjugate is released from the support surface.

- The invention also includes a complex-forming metal ion labeled agent prepared according to the methods of the invention as well as compositions and pharmaceutical compositions including the agent. The invention also includes a composition including a technetium metal labeled agent, wherein the composition includes a specific activity greater than 10,000 Ci/mmol with 99m-technetium and greater than 3,000 Ci/mmol with 188-rhenium. A pharmaceutical composition for radiotherapy or imaging, including a carrier and a complex-forming metal ion labeled agent, wherein the agent is prepared without HPLC.
- 10 Another embodiment of the invention includes a kit for preparing a complex-forming metal ion labeled agent. The kit includes a metal support surface, a conjugate and a predetermined quantity of a complex-forming metal ion. The conjugate is capable of being releasably bound to the support surface and is capable of coordinating with the complex-forming metal ion so that the conjugate is released from the metal support surface.
- 15 The invention also includes a method of detecting the presence or assessing of the severity of a disease, disorder or abnormal physical state in a mammal. The method includes the steps of: (a) administering an effective amount of a compound useful for generating a complex-forming metal ion labeled agent; and (b) detecting the presence or assessing the severity of the disease, disorder or abnormal physical state. Another embodiment includes
- 20 a method of radiotherapy of a disease, disorder or abnormal physical state in a mammal including the steps of administering an effective amount of a compound useful for generating a complex-forming metal ion labeled agent.

- The invention also includes a method for the preparation of a support surface for manufacturing a complex-forming metal labeled agent including electro or electroless metal
- 25 plating or vapor deposition of a suitable thickness, of the metal onto an inorganic or polymeric substrate in the form of particles, sponges or sieves, fibers or surfaces with suitable surface area between about 1 and 10,000cm².

BRIEF DESCRIPTION OF THE FIGURES

Preferred embodiments of the invention will be described in relation to the drawings in which:

5 Figure 1 (a) is a schematic figure showing transchelation of a conjugate (including surface binding group (SBG) and three accessory groups (AG)) from a metal support surface to a complex-forming metal (CFM);

Figure 1(b) a schematic figure showing transchelation of a conjugate from a gold surface to technetium.

Figure 2 is a schematic figure showing *in situ* production of a thiol bound to a gold surface

10 Figure 3 is a radiometric trace of Tc-99m RP895(1) (Labeling of RP414)

Figure 4 is a ultraviolet light trace of Tc-99m RP895(1)

DETAILED DESCRIPTION OF THE INVENTION

2091400 011660

The present invention simplifies the process of generating a metal-ligand complex by coupling a conjugate to a support surface and transchelating the conjugate from the solid support to the metal. In a preferred embodiment, transchelation is from gold to technetium. The support surface preferably includes gold or an element having properties of binding to the metal cleavable conjugate that are similar to gold (gold is preferably Au(111)) so that a metal labeled radiotherapeutic agent or imaging agent can be formed. Metals with properties similar to gold include silver and copper. Preferably, the support surface is at least 50% metal, at least 90% metal and most preferably substantially pure metal (i.e. at least 99.9% metal). The support surface is preferably solid but may be semi-solid (for example, gold may be vapor deposited on a semi-solid compound, such as a soft plastic or resin). The step of introducing a metal to the support surface results not only in the formation of metal-support surface complexes, but also in the consequential release of those complexes from the support for collection in a form substantially free of uncomplexed conjugate. These processes are shown in Figures 1(a) and 1(b). Articles that describe materials and methods for binding molecules to gold include Bain et al., "Formation of

Monolayer Films by the Spontaneous Assembly of Organic Thiols from Solution onto Gold¹", American Chemical Society, 111, 321-335 (1989); Laibinis et al., "Comparison of the Structures and Wetting Properties of Self-Assembled Monolayers of *n*-Alkanethiols on the Coinage Metal Surfaces, Cu, Ag, Au¹", American Chemical Society, 113, 7152-7167 (1991); Sasaki et al., "Two-Dimensional Arrangement of a Functional Protein by Cysteine-Gold Interaction: Enzyme Activity and Characterization of a Protein Monolayer on a Gold Substrate", Biophysical Journal, Volume 72, 1842-1848 (1997); Bain et al., "Correlations between Wettability and Structure in Monolayers of Alkanethiols Adsorbed on Gold¹", American Chemical Society, 110, 3665-3666 (1988) which are incorporated by reference in their entirety.

The present invention also permits the *in situ* formation of the bound thiol through binding to a gold surface. The protected sulfur atom binds to the gold surface resulting in the release of the protecting group as shown in Figure 2.

Compound Aspects

According to an aspect of the present invention, compounds are provided that are useful for generating metal labeled imaging agents that are substantially free of the unlabeled imaging agent. The compounds include a metal support surface, and a conjugate releasably bound to the support surface. The conjugate is capable of coordinating with a complex-forming metal ion so that the conjugate is released (transchelated) from the support surface. The conjugate preferably includes a ligand and a targeting molecule. The conjugate is preferably a peptide, a polypeptide, a peptide or polypeptide mimetic (preferably a derivative) or an organic molecule. Those skilled in the art recognize that a variety of techniques are available for constructing peptide mimetics with the same or similar desired biological activity as a corresponding peptide or polypeptide compound of the invention but with more favorable activity with respect to solubility, stability, and/or susceptibility to hydrolysis and proteolysis. See for example, Morgan and Gainor, *Ann. Rep. Med. Chem.*, 24:243-252 (1989). The organic molecule is preferably a small organic molecule, having molecular weight less than about 600 Daltons and more preferably less than about 500 Daltons. The ligand may be a peptide, a peptide mimetic or a small organic molecule. The ligand preferably incorporates (a) a surface binding group selected from the group consisting of a cysteine amino acid residue, a cysteine amino acid residue derivative, a thiol or thioether group attached to an organic molecule,

- an amino acid residue derivative including phosphorous and a phosphorous containing organic molecule, wherein the amino acid residue, amino acid residue derivative or organic molecule is capable of releasably binding to the support surface; and (b) at least one accessory group capable of coordinating with the complex-forming metal ion. In a variation of the invention it is possible that one or more accessory groups is in the targeting molecule. The conjugate is capable of coordinating with a complex-forming metal ion so that the conjugate is released from the support surface. The conjugate preferably includes a peptide sequence such as a bombesin 7-14 fragment, QWAVGHLM, TKPPR, RGDS, cyclopropylcarbonyl-nLe-L-F-W-E-K(GCSDMG)-G, an antibody (preferably monoclonal or polyclonal) antibody fragment and small organic molecule that targets a receptor or a transporter.

The compound may optionally include a linker, which is, preferably, a biologically inert organic residue connecting the targeting molecule and the conjugate. Suitable linkers are alkyl, polyether, aromatic or polyaromatic compounds, with 3-100 atoms.

- These compounds are useful for generating compositions (preferably imaging agent compositions or radiotherapeutic pharmaceutical compositions) with very high specific activity because they are substantially free of unlabeled agent.

Ligand Definition

- The term "ligand" refers to compounds that incorporate at least one surface binding group. A surface binding group includes a sulfur or phosphorous atom (or atom with similar binding properties) which releasably binds to the support surface. In a preferred embodiment, the atom is sulfur which is capable of binding to gold to become a sulfide. The ligand also includes at least one accessory group, and preferably three of the accessory groups, capable of forming a coordinate bond with a given complex-forming metal, thereby forming a stable metal-ligand complex. The ligand preferably incorporates 3 accessory groups selected from the group consisting of (a) a nitrogen atom, an oxygen atom or a sulfur atom incorporated in an amino acid residue, (b) a nitrogen atom, an oxygen atom, a selenium atom, a phosphorous atom or a sulfur atom incorporated in an amino acid residue derivative, or (c) a nitrogen atom, an oxygen atom, a selenium atom, a phosphorous atom or a sulfur atom incorporated in an organic molecule or (d) a combination of one or more of (a) to (c), wherein the atoms have metal coordinating activity.

In the case where a ligand contains one accessory group and a surface binding group, the ligand may be referred to as a "chelator". Ligands that contain two or more accessory groups and a surface binding group, referred to as multidentate, typically form more stable metal-ligand complexes than do monodentate ligands and are preferred for this reason.

- 5 Many ligands that bind to radionuclide metals are tetradentate containing a combination of four nitrogen and sulfur metal-coordinating atoms i.e. N_3S and N_2S_2 , however they may incorporate other metal-coordinating atoms such as oxygen, phosphorous and selenium (for example see Figure 1(a)). Preferred ligands of the invention include molecules incorporating a sulfur surface binding atom and at least 3 nitrogen atoms (for example see
- 10 Figure 1(b)). The ligand preferably includes a peptide selected from the group consisting of a tetradentate N_xS_{4-x} ligand, a tetradentate N_xS_{4-x} ligand derivative, a polyamino polysulfide and a polyamino polysulfide derivative.

- For radiotherapy and diagnostic imaging it is particularly desirable that the metal complex is highly stable *in vivo* so that the metal is not released from the ligand in substantial quantities
- 15 to accumulate in tissues. The present invention can be applied to a wide variety of ligands, such as N_3S chelators described in PCT application PCT/CA94/00395 and N_2S_2 chelators described in PCT application PCT/CA94/00479. Preferred ligands are peptides or derivatives thereof which incorporate a pendant sulfhydryl group for binding to a metal. Suitable peptidic chelators are those described in WO 9317719 which are amenable to
- 20 coupling to targeting molecules, particularly targeting molecules that are also peptidic. In one embodiment, the invention is applied to label ligands that have intrinsic targeting properties. One such ligand available for radiodiagnostic imaging is mercapto-acetyl-glycyl-glycyl-glycine (MAG3) which localizes in renal tissue and may be labeled according to the method of the present invention to prepare renal imaging or radiotherapeutic agents.
- 25 MAG3 is an N_3S class of ligand having three nitrogen coordinating atoms and one sulfur coordinating atom.

Targeting Molecule

- Targeting molecules suitable for use in compounds of the invention are compounds that are capable of localizing selectively *in vivo* at sites for imaging such as at a particular organ,
- 30 tissue or cell type. Suitable targeting molecules include a polypeptide, a peptide, a nucleic acid molecule, an oligonucleotide, a saccharide, an oligosaccharide, a steroid, a cyclic peptide, a peptide or polypeptide mimetic, an enzyme substrate, an inhibitor and a

small organic molecule. Preferred targeting molecules include polypeptides and peptides, particularly those capable of binding with specificity to cell surface receptors characteristic of a particular pathology. The targeting molecule is preferably a molecule having agonist or antagonist activity. The targeting molecule preferably includes a molecule selected from the group consisting of a bombesin 7-14 fragment, QWAVGHLM, TKPPR, RGDS and small organic molecule that targets a receptor or a transporter. The receptor or transporter is preferably a dopamine receptor or transporter, a serotonin receptor or transporter, a sigma receptor, a GABA receptor, a nicotinic receptor, a cholinergic receptor, a norepinephrine receptor or transporter, a glucose transporter and an opioid receptor. Other preferable targeting molecules are peptides, polypeptides or derivatives comprising 3 or more amino acid residues that bind to cell surface receptors such as those described in PCT/CA94/00395. Preferably, targeting molecules are peptides comprising about: 3 to 1000, 3 to 500, 3 to 100, 3 to 50 amino acids and more preferably 3 to 10 or 3 to 6 amino acids. Small organic molecules of preferably about: 6 to 500, 6 to 250, 6 to 100 carbons and more preferably about 6 to 50 or 6 to 25 carbons are also useful targeting molecules. In an embodiment, targeting molecules are chemotactic peptides that bind to cell surface receptors and in particular are chemotactic peptides that incorporate the amino acid sequence cyclopropylcarbonyl-nLe-L-F-W-E-K(GCSDMG)-G.

In the particular process aspect of the invention, it is desirable that the targeting molecule itself be free of metal binding sites such as pendant sulfhydryl groups. A ligand-targeting molecule conjugate labeled according to this particular process wherein the targeting molecule presents metal binding sites such as pendant sulfhydryl groups found in cysteine residues may 1) lose some or all of its localizing activity and 2) release metal in vivo thereby increasing background noise and obscuring the image.

Ligands and/or targeting molecules that are peptidic are commercially available or may be synthesized de novo by solid phase techniques or by recombinant DNA techniques. Solid-phase peptide synthesis generally involves the use of automated synthesizers and an appropriate support as the solid phase, to which is attached the C-terminal amino acid of the desired peptide. Extension of the peptide in the N-terminal direction is then achieved by successively coupling a suitably protected form of the next desired amino acid, using either Fmoc- or BOC-based chemical protocols typically, until synthesis is complete. Protecting groups are then cleaved from the peptide, usually simultaneously with cleavage of the peptide from the support, and the peptide is then isolated. Common purification techniques

include reversed-phase HPLC using acetonitrile as solvent and trifluoroacetic acid as an ion-pairing agent. Procedures are described in numerous publications. Reference may be made to Stewart and Young, Solid Phase Peptide Synthesis, 2nd Edition (and subsequent editions), 1984, Pierce Chemical Company, Rockford, Illinois which is incorporated by
5 reference in its entirety. Alternatively, peptides may be synthesized in solution or on a solid phase in small blocks and subsequently ligated to give the desired sequence. Peptides incorporating amino acids that are not genetically encoded require synthetic techniques for preparation.

Support

10 The term "support surface" refers to any substrate that is insoluble and inert in labeling solutions and is a metal support surface which is either made of, or is coated with, gold, silver or copper or a metal capable of releasably binding and coordinating sulfur or phosphorous for forming a metal complex in preparation of labeled agents. Suitable
15 compounds that may be coated with a metal include inorganic silicate glass, alkylamino functionalized controlled-pore glass, silica or alumina beads and organic polystyrene, polyacrylamide or sugar polymers such as Sephadex and agarose. The support surface may be provided as a powder, a solid piece of metal, a ball contained in a tube or as a coating on the inside of a vessel. For the purposes of generating metal-ligand complexes the metal support may be in a column that allows for easy passage, collection
20 and filtration of the complex solution.

The invention also includes a method for the preparation of a support surface for manufacturing a complex-forming metal labeled agent comprising electro or electroless metal plating or vapor deposition of a suitable thickness of the metal onto a preferably inorganic or polymeric substrate in the form of particles, sponges or sieves, fibers or surfaces
25 with suitable surface area preferable between about 1 and 10,000cm². The thickness of the metal on the metal support surface is preferably greater than about 10nm.

Metal

The term "complex-forming metal" refers to any metal atom that is in a state capable of forming stable coordination bonds with metal coordinating atoms of a ligand. Complex-
30 forming metals include the transition metals, lanthanide metals and actinide metals.

Complex-forming metals useful in preparing agents for radiotherapy or imaging are preferably the metals (or radioisotopes of the metals) Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.

5 For MRI, the complex-forming metal can be a paramagnetic metal atom such as divalent and trivalent chromium, manganese, iron, cobalt, nickel, copper, samarium,. The more preferred complex-forming metals for MRI are those exhibiting a strong magnetic moment, such as manganese.

10 The halide salt, in particular chloride salt, or oxide of these complex-forming metals are forms capable of complexing with a desired ligand and are suitable for the present invention. Radionuclide labeled imaging agents employ complex-forming metal isotopes that include β -emitters such as rhenium-186 and -188; and γ -emitters such as technetium-99m. The complex-forming metal most preferred for radiodiagnostic imaging is technetium-99m due to its advantageous half life of 6 hours and inexpensive preparation from a molybdenum-99 generator. Technetium and rhenium labeling is accomplished by
15 procedures established in the art. Either complex-forming metal may be introduced to the ligand in aqueous solution in oxo, dioxo or nitrido form, for example pertechnetate ($^{99m}\text{TcO}_4^-$) or perrhenate, with a suitable reducing agent such as stannous chloride. Technetium may also be reduced by electrochemical reduction. Alternatively, radiodiagnostic agents may be formed by a transchelation reaction which entails use of the complex-forming metal in the
20 form of a weak metal complex such as technetium-gluconate, heptagluconate, tartrate or citrate to give a desired labeled ligand. Transchelation reactions are typically heated to facilitate conversion of technetium from the weak complex to a complex with the ligand, for example in a boiling hot water bath.

Method Aspects

25 According to a method of the present invention, ligand-targeting molecule conjugates are labeled with complex-forming metals to provide a solution substantially free from unlabeled conjugate. The method for generating a complex-forming metal labeled conjugate diagnostic agent or radiotherapeutic agent includes transchelating a conjugate from a metal support surface to a complex-forming metal ion so that the conjugate is released from
30 the support surface. In general, the process comprises the steps of obtaining a compound in which the conjugate is coupled to a support surface via a metal-cleavable surface binding

group; introducing a complex-forming metal to the support; and collecting the labeled conjugate released from the support. In this process the complex-forming metal forms a coordination bond with the surface binding group and the accessory group(s) of the ligand thereby cleaving the bond between the ligand atom and the support. As a result, only

5 labeled conjugate is released from the support. The complex-forming metal labeled agent has very high specific activity > about 10,000 Ci/mmol with 99m-technetium and > about 3,000 Ci/mmol with 188-rhenium.

In one preferred embodiment, the method for generating a complex-forming metal labeled conjugate diagnostic agent or radiotherapeutic agent, includes (a) obtaining a suitable

10 compound prepared as described in this application and (b) contacting the compound with the complex-forming metal ion to form a coordinate bond between the complex-forming metal ion and the agent so that the complex-forming metal labeled agent is released from the support surface. Preferably, the method includes collecting the complex-forming metal labeled agent so released.

15 In another embodiment, the gold surface is employed for *in situ* formation of the bound thiol. This eliminates several steps which are detrimental to subsequent kit formulation. These steps cause difficulties in ensuring sterility.

A preferred method for preparing a complex-forming metal ion labeled agent (eg. peptide or polypeptide) includes (a) bonding a peptide, polypeptide, peptide mimetic or polypeptide

20 mimetic that includes (i) a cysteine amino acid residue, a cysteine amino acid residue derivative or a phosphorous amino acid residue derivative and (ii) at least one accessory group capable of coordinating with the complex-forming metal ion, to a metal support surface by the sulfur atom or phosphorous atom of the residue or derivative and (b) labeling the peptide, polypeptide or mimetic with a complex-forming metal so that the metal

25 complexed peptide, polypeptide or mimetic is released from the support surface. The peptide, polypeptide, peptide mimetic or polypeptide mimetic is preferably between about 3 and 50 amino acid residues or amino acid residue derivatives.

- A preferred method for preparing a complex-forming metal ion labeled organic molecule includes (a) bonding an organic molecule that includes (i) a sulfur atom or a phosphorous atom and (ii) at least one accessory group capable of coordinating with the complex-forming metal ion, to a support surface by the sulfur group or the phosphorous atom and
- 5 (b) labeling the agent with a complex-forming metal so that the metal complex conjugate is released from the support surface. The organic molecule preferably includes between about 6 and 100 carbons.

- The preferred support surfaces are as described above and include metals such as gold, silver, copper and a metal capable of releasably binding sulfur or phosphorous for forming a
- 10 metal ion labeled agent. The complex-forming metal is preferably a metal or radioisotopic form of metal from the group of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.

The invention also includes the complex-forming metal ion labeled agent product prepared according to a method of the invention.

15 **Pharmaceutical compositions**

- The invention also includes compositions, preferably pharmaceutical compositions for radiotherapy or imaging, including an agent prepared according to a method of the invention. The agents and compositions are preferably prepared without HPLC. The invention provides a composition including an agent (prepared according to a method of the
- 20 invention, or any other method) which includes 99m-technetium metal labeled agent (preferably prepared on a gold support surface), and which has a specific activity greater than about 10,000 Ci/mmol with 99m-technetium or greater than about 3,000 Ci/mmol with 188-rhenium. The agent is preferably a peptide such as
- 25 dimethylglycylserinylcysteinylglycine or polypeptide or a mimetic (preferably a derivative) thereof,. Pharmaceutical compositions may be formulated according to known techniques.

- The invention includes a method of detecting the presence or assessing of the severity of a disease, disorder or abnormal physical state in a mammal comprising: (a) administering an agent or composition of the invention and (b) detecting the presence or assessing the severity of the disease, disorder or abnormal physical state. The presence or severity of
- 30 the disease, disorder or abnormal physical state is detected or assessed with a technique

selected from the group consisting of positron emission tomography, nuclear magnetic resonance imaging, scintigraphy, single photon emission computed tomography, perfusion contrast echocardiography, ultrafast X-ray computed tomography, and digital subtraction angiography. Preferably, the agent includes a ^{99m}Tc metal and binds to a receptor and the technique is single photon emission computed tomography. Suitable methods and materials for imaging are described in: Handbook of Nuclear Medicine second ed., 1993, Mosby Press, Frederic I. Datz; Fundamentals of Nuclear Pharmacy third ed., 1992 Springer-Verlag, Gopal B. Saha; Principles and practice of Nuclear Medicine second ed., 1995, Mosby press, Paul J. Early and D. Bruce Sodee; which are incorporated by reference in their entirety.

The invention also includes a method of radiotherapy of a disease, disorder or abnormal physical state in a mammal including administering an agent or composition of the invention. Methods of performing radiotherapy are described in, for example, Principles and Practice of Nuclear Medicine, 2nd Ed., P.J. Early and D.B. Sodee, Chapter 32, which is incorporated by reference in its entirety.

The pharmaceutical compositions are used to treat diseases and provide images in diseases, disorders or abnormal physical states including oncological, neurological, inflammatory, infection, and degenerative diseases. Other diseases, disorders and abnormal physical states will be apparent to those skilled in the art and/or on review of this application or references cited in this application.

Pharmaceutical compositions used for imaging or to treat patients having diseases, disorders or abnormal physical states preferably include an agent of the invention and an acceptable vehicle or excipient (Remington's Pharmaceutical Sciences 18th ed, (1990, Mack Publishing Company) and subsequent editions). Vehicles include saline and D5W (5% dextrose and water). Excipients include additives such as a buffer, solubilizer, suspending agent, emulsifying agent, viscosity controlling agent, lactose filler, antioxidant, preservative. The compositions may further include a reducing agent, a bulking agent or a pH stabilising agent. There are preferred excipients for stabilizing peptides for parenteral and other administration. The complex-forming metal labeled agent is preferably administered by an intravenous parenteral route. The excipients preferably include serum albumin, glutamic or aspartic acid, phospholipids and fatty acids. Parenteral (injectable) administration is preferred. The methods for the

preparation of pharmaceutically acceptable compositions which can be administered to patients are known in the art.

The pharmaceutical compositions can be administered to humans or animals (preferably mammals). Dosages to be administered depend on individual patient condition,

- 5 indication of the drug, physical and chemical stability of the drug, toxicity, the desired effect and on the chosen route of administration (Robert Rakel, ed., Conn's Current Therapy (1995, W.B. Saunders Company, USA)). Preferably the amount of complex-forming metal labeled agent administered to the mammal is about 0.01 mcg/kg/minute to 1,000 mcg/kg/minute and more preferably about 0.01 to 50 mcg/kg/minutes.

10 Kits

The invention also includes kits for preparing a complex-forming metal ion labeled agent.

The kit preferably includes a metal support surface, a conjugate and a predetermined quantity of a complex-forming metal ion, the conjugate being capable of being releasably bound to the support surface and capable of coordinating with the complex-forming

- 15 metal ion so that the conjugate is released from the metal support surface. Suitable support surfaces, conjugates and complex-forming metal ions are described in this application or apparent to a person of skill in the art upon reviewing this application. The kit preferably includes a conjugate including a ligand and a targeting molecule. The ligand preferably incorporates (a) a surface binding group selected from the group consisting of
- 20 a cysteine amino acid residue, a cysteine amino acid residue derivative, an organic thiol or thioether containing molecule, an amino acid residue derivative including phosphorous and a phosphorous containing aliphatic molecule, wherein the amino acid residue, amino acid residue derivative or aliphatic molecule is capable of releasably binding to the support surface; and (b) at least one accessory group. The conjugate is capable of
- 25 coordinating with a complex-forming metal so that the conjugate is released from the support surface. Preferable targeting molecules are described in this application. The polypeptides, peptides or other reagents may be lyophilized. Since the radioisotopic metals often have a very short half life, it is advantageous to omit them from the kit.

- 30 The support surface preferably includes a metal selected from the group consisting of gold, silver, copper and a metal capable of releasably binding sulfur or phosphorous for forming a metal complex. The complex-forming metal is preferably one or more of the

metals and radioisotopic metal forms of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.

Preferred embodiments of the invention are described below in examples which are not intended to in any way limit the scope of the invention.

5 **Example 1 - Preparation of dimethylglycylserinylcysteinylglycine**

A solution of dimethylglycylserinylcysteinyl(Acetomidomethyl)glycine (RP414) (50mg) and mercuric acetate (75mg) in 30% acetic acid (2mL) was stirred at room temperature for 3h. Distilled water (2mL) was added and H₂S gas was slowly bubbled through the solution for 3 min causing a black precipitate (mercury sulfide). The solution was filtered
10 through a 0.2 micron filter diluted with a further 2mL of distilled water and lyophilised.

Example 2 - Preparation of dimethylglycylserineylcysteinylglycine (RP414-SH) attached to a gold surface.

A piece of gold foil (50 micron thickness) measuring 1 x 5cm was coiled into a spiral and placed into a test tube. To this spiral was added a solution of 20% nitric acid and the
15 whole gently agitated for 2h. The nitric acid was removed and the foil washed with distilled water (5 times 4mL for 30 min each). The final portion of water was removed and the foil soaked in phosphate buffered saline (0.01M) for 1.5h at pH 7. The phosphate was then removed immediately prior to addition of the thiol.

A solution of dimethylglycylserinylcysteinylglycine (RP414-SH) prepared as in example 1
20 (~50mg) in phosphate buffered saline (3mL) was adjusted to pH 7 using KHCO₃ / K₂CO₃ buffer. This solution was added to the foil prepared above and the reaction mixture gently agitated at room temperature for 20h under an argon atmosphere. The solvent was decanted off and the foil washed with successive portions (2mL) of phosphate buffered saline (five portions for 30min each) with gentle agitation. The final aliquot was
25 retained for analysis by HPLC to ensure that all non-adsorbed RP414-SH was removed. The final aliquot was decanted off and the apparatus purged with argon ready for labelling with 99m-Tc. The peptide RP414-SH adsorbed to the gold surface was designated RP895.

Radiolabelling Method of Gold Surface

- 1.0mL of saline was added to the above prepared RP895
- 100µL of stannous gluconate was added (stannous gluconate is prepared by the addition of 20µL of 20mg/mL stannous chloride to 1.0mL of 13mg/mL sodium gluconate)
- approximately 10mCi of Tc-99m pertechnetate in 200µL of saline was added to the reaction
- The reaction vessel was then shaken vigorously
- Allow the reaction to take place for 1 hour at room temperature
- Then incubate reaction for 30 minutes at 60°C

The total reaction volume was 1.3mL, a clear and colourless solution with a pH of 5.0 was achieved.

- A 500µL sample was analyzed using high performance liquid chromatography with a reverse phase C-18 column and a step gradient method of 0 to 70% acetonitrile w/0.1% TFA over 25 minutes. The sample was analyzed by both radiometric and UV detectors (see Figure 3 for radiometric and Figure 4 for UV of the final reaction.

Example 3 - Radiolabelling Method for RP414 on Gold Flakes (RP902)

- 15 – 830 mg of gold flakes loaded with RP414 was soaked in 1.0 ml of saline, followed by the addition of 0.1 ml of stannous gluconate and 0.1 ml of Tc-99m sodium pertechnetate (approximately 10 mCi). The reaction which contains a small stirrer was shaken vigorously and placed on a hot plate/stirrer to be incubated at 60 – 70°C for 1 hour. An aliquot of the reaction mixture was removed and filtered through an acrodisc low protein binding 0.2µm membrane filter. The filtered reaction was then analyzed using HPLC with a reverse phase C18 column and a step gradient of 0.1% TFA in both water and acetonitrile. The sample was analyzed by both radiometric and UV detectors.

Example 4 - Preparation of dimethylglycine-serine-cysteine-glycine on solid support

RP414 (dimethylglycine-serine-cysteine (Acm)-glycine) 50mg was dissolved in 30% acetic acid and mercury acetate (75mg) added. The mixture was stirred at room temperature for 3.5h hydrogen sulfide gas was bubbled through the solution with the formation of a black precipitate. This precipitate was removed by filtration and the solvents removed from the filtrate under reduced pressure.

Gold powder (2 micron average particle size) (75mg) was wetted a drop of methanol in distilled water and the solution removed. The powder was then suspended in 20% nitric acid and shaken at 50C for 1.5h and after settling the supernatant removed. The powder was then washed with distilled water (4 times 5mL for 30 min each) and then washed with phosphate buffered saline (2mL). After removal of the saline a solution of de-protected RP414 in phosphate buffered saline (2mL) was added and the mixture shaken at room temperature under argon for 20h. The supernatant was removed and the gold powder washed with successive portions of phosphate buffered saline (5 times 4mL).

Example 5: Attachment of RP527-SH to Gold Flakes/Spheres (RP904)

Gold powder (1.5-3.0 micron particles, ~100 mg) was flamed using a bunsen burner. Methanol was added to the gold and stirred vigorously to break up the gold into fine powder. The suspension was centrifuged and the solution decanted.

RP527(dimethylglycine-ser-cys(Acm)-gly- β ala-gln-trp-ala-val-gly-his-leu-met-NH₂) (~10mg, 1eq) was dissolved in 30% acetic acid (4mL) and then mercuric acetate (2.2 mg) was added. The reaction was left at room temperature for 3 hours and then bubbled through with hydrogen sulfide for 2 minutes. The resulting mixture was centrifuged and the solution decanted from the pellet of mercury sulfide. The solvent was removed in vacuo. RP527-SH (~10 mg) was then dissolved in a 1:1 solution of ethanol:0.01 PBS and added to the gold powder in a vacutainer with vigorous stirring (magnetic stirbar was added). The solution was flushed with argon for 10 minutes and left under argon at room temperature for 24 hours. The reaction mixture was centrifuged and the supernatant removed. The powder was washed with a 1:1 mixture of ethanol:aqueous 0.1% trifluoroacetic acid (5 x 4 mL), centrifuging between each addition of washing solution.

The final aliquot was decanted off and the gold was dried in vacuo. It was purged with argon and stored at -20°C in preparation for labeling with Tc-99m.

5 **Radiolabelling Method for RP527(dimethyglycine-ser-cys(Acm)-gly-βala-gln-trp-ala-val-gly-his-leu-met-NH₂) on Gold Flakes (RP904)**

100 mg of gold flakes loaded with RP527 was soaked in 1.0 ml of saline, followed by the addition of 0.1 ml of stannous gluconate and 0.1 ml of Tc-99m sodium pertechnetate (approximately 10 mCi). The reaction which contains a small stirrer was shaken vigorously and placed on a hot plate/stirrer to be incubated at 70 - 75°C for 1 hour. An aliquot of the reaction mixture was removed and filtered through an acrodisc low protein binding 0.2µm membrane filter. The filtered reaction was then analyzed using HPLC with a reverse phase C18 column and a step gradient of 0.1% TFA in both water and acetonitrile. The sample was analyzed by both radiometric and UV detectors.

15 **Example 6: Preparation of RP414 (dmG-S-C(Acm)-G) attached to gold foil (RP895)**

A piece of gold foil (50 micron thickness, 1 cm x 5 cm) was folded into an accordion shape and subsequently shaped into a cylinder. A solution of 20% nitric acid (4 mL) was added to the gold foil and heated at 40°C for 2h. The nitric acid was removed and the foil was washed with distilled water (5 x 4 mL) for a few minutes. RP414 (~50 mg) was dissolved in 30% acetic acid (4 mL) and added to the gold in a vac tube. The solution was flushed with argon and left under argon at room temperature for 24 hours. The supernatant was removed and the foil washed with distilled water (5 x 4 mL). The final aliquot was decanted off and the vac tube was purged with argon in preparation for labeling with Tc-99m.

This experiment was repeated with RP128 (dmG-S-C(Acm)-G-T-K-P-P-R) attached to gold foil (RP905).

30 **Example 7: Preparation of Kit Formulation for RP895**

A piece of gold foil (50 micron thickness, 1cm x 5 cm) was prepared and washed using the same procedure outlined in Example 1. RP414 (~50 mg) was loaded onto the gold using the same procedure outlined in Example 1. Once washed, labeling solution

consisting of stannous gluconate (100 μ L), stannous chloride (20 μ L, 20mg/mL) and sodium gluconate (1mL, 13mg/mL) was added, frozen and lyophilized.

Example 8: Preparation of RP414 attached to gold powder (RP902)

5

Gold powder (1.5-3.0 micron particles, ~75 mg) was wetted with 5% methanol in water. The suspension was centrifuged and the solution decanted. The gold powder was washed with water (4 mL) and centrifuged. The water was decanted and 20% nitric acid was added to the gold. The suspension was heated at 40°C for 2h. The suspension was centrifuged and the nitric acid decanted. The powder was washed with distilled water (5 x 4 mL), each time centrifuging between each addition of water. RP414 (~50 mg) was dissolved in 30% acetic acid (4 mL) and added to the gold powder in a vac tube with vigorous stirring (magnetic stirbar was added). The solution was flushed with argon and left under argon at room temperature for 24 hours. The supernatant was removed and the powder washed with distilled water (5 x 4 mL), centrifuging between each addition of water. The final aliquot was decanted off and the vac tube was purged with argon in preparation for labeling with Tc-99m.

20

Example 8a:

RP902 and RP904 were prepared as described in Example 8. However, the solution in which they were loaded onto the gold powder was changed to 1:1 ethanol:30% acetic acid. The final samples were lyophilized in preparation for labeling with Tc-99m.

25

Example 8b:

Gold powder (~75mg) was loaded with de-protected RP414 in 1:1 ethanol:0.01M PBS. The rest of the reaction was carried out following the procedure in Example 8. This reaction was repeated for de-protected RP527.

30

Example 9: Incremental Labeling of RP414 on Gold Powder (RP902)

A large batch of gold powder (1.5-3.0 micron particles, ~2g) was washed and loaded with RP414 as described in Example 3. Ever increasing amounts were prepared for labeling

35

with Tc-99m to investigate the efficiency of labeling as the amount of RP414 on gold powder was increased. Amino acid analysis was carried out on this batch of RP902.

Amounts of RP902 submitted for Tc-99m labeling:

5

7.85mg, 17.9mg, 34.4mg, 78.8mg, 151.3mg and 308.6mg

EXAMPLE 10: METHODS OF RADIOLABELLING GOLD COMPOUNDS WITH Tc-99m PERTECHNETATE

10 Preparation of Stannous Gluconate:

0.02mL of 20mg/mL stannous chloride was added to 1.0mL of 13mg/mL sodium gluconate and shaken well

1. Radiolabelling Method for RP128 on Gold Foil (RP905)

15 A piece of RP905 was soaked with 1.5mL of saline in a glass vacutainer. 0.1mL of freshly prepared stannous gluconate was added as well as 0.1mL of Tc-99m pertechnetate (approximately 10mCi). The reaction vessel was shaken vigorously and allowed to sit at room temperature for 1 hour. The final reaction volume was 1.7mL and was clear and colourless with a pH of 4.5 – 5.5. An aliquot of the reaction was analyzed
20 using HPLC with a reverse phase C-18 column and a step gradient method of 0 to 70% acetonitrile w/ 0.1% TFA over 25 minutes. The sample was analyzed by both radiometric and UV detectors.

The same method was repeated with RP414 on gold foil (RP895).

25

2. Radiolabelling Method for RP527 on Gold Foil (RP900)

A piece of RP900 was soaked with 1.0mL saline in a glass vacutainer, followed by the addition of 0.1mL of stannous gluconate and 1.0mL of Tc-99m pertechnetate (approximately 10mCi). The reaction vessel was shaken vigorously and placed in a
30 water bath for incubation at 55°C for 1 hour. The final reaction solution was clear and colourless with a pH of 4.5 – 5.0. An aliquot of the reaction is analyzed using HPLC with a reverse phase C-18 column and a step gradient method of 0 to 70% acetonitrile w/ 0.1% TFA over 25 minutes. The sample was analyzed by both radiometric and UV detectors (wavelength = 214nm).

35

3. Radiolabelling Method for RP414 on Gold Powder (RP902)

75mg of gold powder was loaded with RP414 was soaked in 0.5mL of saline, followed by the addition of 0.1mL of stannous gluconate and 0.1mL of Tc-99m pertechnetate (approximately 10mCi). The reaction was then shaken vigorously and stirred in an oil bath at 55 - 60°C for 1 hour. Care was taken when removing the reaction out of the water bath so as not to disturb the powder, which settled to the bottom of the container. This portion was then filtered through a 0.2 μ acrodisc filter to give a clear solution with a pH between 4.5 - 5.0. An aliquot of the reaction was analyzed using HPLC with a reverse phase C-18 column and a step gradient method of 0 to 70% acetonitrile w/ 0.1% TFA over 25 minutes. The sample was analyzed by both radiometric and UV detectors (wavelength = 214nm).

The same method was used for RP527 except the temperature and time of incubation was changed to 100°C for 35 minutes.

The present invention has been described in detail and with particular reference to the preferred embodiments; however, it will be understood by one having ordinary skill in the art that changes can be made without departing from the spirit and scope thereof. For example, where the application refers to peptides, it is clear that polypeptides and proteins may often be used.

All publications, patents and patent applications are incorporated by reference in their entirety to the same extent as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety.

WE CLAIM

1. A composition for generating a complex-forming metal ion labeled agent, the composition comprising:

- (a) a metal support surface; and
- (b) a conjugate releasably bound to the support surface, the conjugate comprising a ligand and a targeting molecule;

wherein the conjugate is capable of coordinating with a complex-forming metal ion so that the labeled conjugate is released from the support surface.

2. The composition of claim 1, wherein the metal support surface is capable of releasably coordinating to sulfur or phosphorous and the ligand comprises a sulfur or phosphorous atom for binding to the metal support surface.

3. The composition of claim 2, wherein the ligand comprises a sulfur atom attached to a sulfur protecting group, the metal support surface being capable of binding to the protected sulfur atom thereby releasing the sulfur protecting group from the sulfur atom and forming a thiol bond with the ligand.

4. The composition of claim 2, wherein the conjugate comprises a peptide, a polypeptide, a peptide or polypeptide mimetic or a small organic molecule.

5. The composition of claim 4, wherein the conjugate comprises a peptide sequence selected from the group consisting of a bombesin 7-14 fragment, QWAVGHLM, TKPPR and RGDS.

6. The composition of claim 4, wherein the conjugate comprises a small organic molecule that targets a receptor or a transporter.

7. The composition of claim 2, wherein the ligand comprises:

- (a) a surface binding group selected from the group consisting of a cysteine amino acid residue, a cysteine amino acid residue derivative, a thiol or thioester group attached to an organic molecule, an amino acid residue derivative including phosphorous and a phosphorous containing organic molecule, wherein the amino acid residue, amino acid residue derivative or organic molecule is capable of releasably binding to the support surface; and
- (b) at least one accessory group capable of coordinating with the complex-forming metal ion.

8. The composition of claim 7, wherein the ligand comprises a peptide, a peptide mimetic, a polypeptide, a polypeptide mimetic or a small organic molecule.

9. The composition of claim 8, wherein the ligand comprises a peptide selected from the group consisting of a tetradentate N_4S_{4-x} ligand, a tetradentate N_4S_{4-x} ligand derivative, a polyamino polysulfide and a polyamino polysulfide derivative.
10. The composition of any one of claims 7 to 9, wherein the ligand comprises 3 accessory groups selected from the group consisting of (a) a nitrogen, oxygen or sulfur atom incorporated in an amino acid residue; (b) a nitrogen, oxygen, selenium, phosphorous or sulfur atom incorporated in an amino acid residue; (c) a nitrogen, oxygen, selenium, phosphorous or sulfur atom incorporated in an organic molecule; and (d) a combination of one or more of (a) to (c), wherein the residues, derivatives and/or molecules have metal coordinating activity.
11. The composition of any one of claims 1 to 10, wherein the targeting molecule comprises a molecule having agonist or antagonist activity selected from the group consisting of a polypeptide, a peptide, a nucleic acid molecule, an oligonucleotide, a saccharide, an oligosaccharide, a steroid, a cyclic peptide, a peptide or polypeptide mimetic, an enzyme substrate, an inhibitor and a small organic molecule.
12. The composition of any one of claims 1 to 11, wherein the targeting molecule comprises a peptide, a polypeptide, a peptide or polypeptide mimetic or a small organic molecule.
13. The composition of any one of claims 1 to 12, wherein the targeting molecule comprises a molecule selected from the group consisting of a bombesin 7-14 fragment, QWAVGHLM, TKPPR, RGDS and a small organic molecule that targets a receptor or a transporter.
14. The composition of either of claims 6 or 13, wherein the receptor or transporter is selected from the group consisting of a dopamine receptor or transporter, a serotonin receptor or transporter, a sigma receptor, a GABA receptor, a nicotinic receptor, a cholinergic receptor, a norepinephrine receptor or transporter, a glucose transporter and an opioid receptor.
15. The composition of any one of claims 1 to 14, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal capable of releasably binding sulfur or phosphorous for forming a metal complex.
16. The composition of any one of claims 1 to 14, wherein the metal support surface comprises gold.
17. The composition of any one of claims 1 to 16, wherein the complex-forming metal is selected from the group of metals and radioisotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.

18. The composition of any one of claims 1 to 16, wherein the complex-forming metal is selected from the group of metals and radioisotopic metals consisting of Tc, Cu and Re.
19. A method for generating a complex-forming metal ion labeled diagnostic agent or radiotherapeutic agent, comprising: (a) providing a composition according to any one of claims 1 to 18; and (b) contacting the composition with the complex-forming metal ion to form a coordinate bond between the complex-forming metal ion and the agent so that the complex-forming metal labeled agent is released from the support surface.
20. The method of claim 19, further comprising collecting the complex-forming metal labeled agent so released.
21. A metal ion labeled agent prepared using a composition of any one of claims 1 to 18.
22. A technetium or rhenium labeled agent prepared using a composition of any one of claims 1 to 18, wherein the agent is labeled with ^{99m}Tc and has a specific activity of greater than 10,000 Ci/mmol or the agent is labeled with ^{188}Re and has a specific activity of greater than 3,000 Ci/mmol.
23. The composition of claim 22, wherein the agent is a peptide comprising dimethylglycylserinylcysteinylglycine.
24. A pharmaceutical composition for radiotherapy or imaging, comprising a carrier and a complex-forming metal ion labeled agent, wherein the agent is prepared using a composition of any one of claims 1 to 18.
25. The pharmaceutical composition of claim 24 further comprising at least one agent selected from the group consisting of a reducing agent, a bulking agent and a pH stabilizing agent.
26. A method of detecting the presence or assessing the severity of a disease, disorder or abnormal physical state in a mammal comprising:
- (a) administering an effective amount of the agent or composition of any one of claims 21 to 25; and
 - (b) detecting the presence or assessing the severity of the disease, disorder or abnormal physical state.
27. A method of radiotherapy of a disease, disorder or abnormal physical state in a mammal comprising administering an effective amount of the agent or composition of any one of claims 21 to 25.
28. The method of either of claims 26 or 27, wherein the complex-forming metal labeled imaging agent is administered by an intravenous route.

29. The method of any one of claims 26 to 28, wherein the amount of complex-forming metal labeled agent administered to the mammal is about 0.01 mcg/kg/minute to 1,000 mcg/kg/minute.
30. The method of claim 29, wherein the amount of the complex-forming metal labeled agent administered to the mammal is about 0.01 to 50 mcg/kg/minute.
31. The method of any one of claims 26 to 30, wherein the mammal is a human.
32. The method of any one of claims 26 to 31, wherein the disease, disorder or abnormal physical state is selected from the group consisting of oncological, neurological, inflammatory, infection, and degenerative diseases, disorders and abnormal physical states.
33. The method of claim 32, wherein the presence or the severity of a disease, disorder or abnormal physical state is detected or assessed with a technique selected from the group consisting of positron emission tomography, nuclear magnetic resonance imaging, scintigraphy, single photon emission computed tomography, perfusion contrast echocardiography, ultrafast X-ray computed tomography, and digital subtraction angiography.
34. The method of claim 33, wherein the agent comprises a ^{99m}Tc metal and binds to a receptor and the technique is single photon emission computed tomography.
35. A kit for preparing a complex-forming metal ion labeled agent, the kit comprising a metal support surface, a conjugate and a predetermined quantity of complex-forming metal ion, the conjugate being capable of being releasably bound to the support surface and capable of coordinating with the complex-forming metal ion so that the conjugate is released from the metal support surface.
36. The kit of claim 35, wherein the conjugate comprises a sulfur atom attached to a sulfur protecting group, the metal support surface being capable of binding to the protected sulfur atom thereby releasing the sulfur protecting group from the sulfur atom and forming a thiol bond with the conjugate.
37. The kit of claim 35, wherein the metal support surface is capable of releasably coordinating to sulfur or phosphorous and the conjugate comprises a sulfur or phosphorous atom for binding to the metal support surface.
38. The kit of claim 37, wherein the conjugate comprises a ligand and a targeting molecule, wherein the ligand comprises:
- (a) a surface binding group selected from the group consisting of a cysteine amino acid residue, a cysteine amino acid residue derivative, a thiol or thioester group attached to an organic molecule, an amino acid residue derivative including phosphorous and a phosphorous containing organic molecule, wherein the amino acid residue, amino acid residue derivative

or organic molecule is capable of releasably binding to the support surface; and

- (b) at least one accessory group capable of coordinating with the complex-forming metal ion.

39. The kit of any one of claims 35 to 38, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal capable of releasably binding sulfur or phosphorous for forming a metal complex; and the complex-forming metal is selected from the group of metals and radioisotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.

40. The kit of any one of claims 35 to 39, further comprising at least one agent selected from the group consisting of a reducing agent, a bulking agent and a pH stabilising agent.

41. A method for generating a complex-forming metal ion labeled agent comprising:

- (a) providing a metal support surface;
- (b) providing a conjugate comprising a ligand and a targeting molecule, wherein the ligand comprises a peptide, a peptide mimetic, a polypeptide or a polypeptide mimetic of about 3 to 50 amino acid residues or derivatives thereof and includes a sulfur atom for binding to the metal support surface, the sulfur atom being protected by a sulfur protecting group;
- (c) contacting the protected sulfur atom with the metal support surface so that the sulfur atom forms a thiol bond with the metal surface thereby releasing the sulfur protecting group; and
- (d) contacting the ligand with the complex-forming metal ion to form a coordinate bond between the complex-forming metal ion and the ligand so that the complex-forming metal labeled agent is released from the support surface.

42. The method of claim 41, wherein the metal support surface comprises a metal selected from the group consisting of gold, silver, copper and a metal capable of releasably binding sulfur for forming a metal complex; and the complex-forming metal is selected from the group of metals and radioisotopic metals consisting of Tc, Re, Mn, Fe, Co, Ni, Zn, Cd, Mo, W, Cu, Ag, Au, Ti, Hg, Cr and Rh.

43. A method for the preparation of a support surface for manufacturing a complex-forming metal labeled agent comprising electro or electroless metal plating or vapor deposition of a suitable thickness of the metal onto an inorganic or polymeric substrate in the form of particles, sponges or sieves, fibers or surfaces with suitable surface area between about 1 and 10,000 cm².

44. The method of claim 43, wherein the thickness of the metal on the support surface is greater than about 10 nm.

w178158v2





Fig.1(a)






Figure 3




Figure 4

5/5



PATENT DECLARATION
FILE NO. 7126-2
DE&Y File No.:2274-020133
PAGE 1 OF 3

DECLARATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or the below named inventors are the original, first and joint inventors (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled **IMMOBILIZED LABELLING COMPOUNDS AND METHODS**, the Specification of which:

- ☐ Is attached hereto.
☒ was filed on August 10, 2001 as Application Serial No. 09/913401.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims.

I acknowledge the duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability of the subject matter claimed in this application, as "materiality" is defined in Title 37, Code of Federal Regulations, § 1.56.

I hereby claim priority benefits under Title 35, United States Code, § 119 and/or § 365 of any foreign application(s) for patent or inventor's certificate, PCT international application(s), and United States provisional application(s), listed below and have also identified below any foreign application for patent or inventor's certificate, PCT international application, or United States provisional application, having a filing date before that of the application on which priority is claimed:

PRIORITY APPLICATION(S)

			Priority Claimed
60/120,561	UNITED STATES	17 February 1999	Yes
(Number)	(Country)	(Date Filed)	Yes/No
PCT/CA99/00700	CANADA	3 August 1999	Yes
(Number)	(Country)	(Date Filed)	Yes/No

I hereby direct that all correspondence and telephone calls be addressed to Barbara S. Kitchell, Akerman, Senterfitt & Eidson, P.A., Fourth Floor, 222 Lakeview Avenue, P.O. Box 3188, West Palm Beach, FL 33402-3188, (561) 653-5123.

I HEREBY DECLARE THAT ALL STATEMENTS MADE OF MY OWN KNOWLEDGE ARE TRUE AND THAT ALL STATEMENTS MADE ON INFORMATION AND BELIEF ARE BELIEVED TO BE TRUE; AND FURTHER THAT THESE STATEMENTS WERE MADE WITH THE KNOWLEDGE THAT WILLFUL FALSE STATEMENTS AND THE LIKE SO MADE ARE PUNISHABLE BY FINE OR IMPRISONMENT, OR BOTH UNDER SECTION 1001 OF TITLE 18 OF THE UNITED STATES CODE AND THAT SUCH WILLFUL FALSE STATEMENTS MAY JEOPARDIZE THE VALIDITY OF THE APPLICATION OR ANY PATENT ISSUED THEREON.




FULL NAME OF INVENTOR	FAMILY NAME POLLAK	FIRST GIVEN NAME ALFRED	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY	STATE OR COUNTRY SLOVENIA	COUNTRY OF CITIZENSHIP SLOVENIA
POST OFFICE ADDRESS	POST OFFICE ADDRESS MURNIKOVA 8A, 1000 LUJBANA	CITY	STATE & ZIP CODE/COUNTRY SLOVENIA
INVENTOR'S SIGNATURE:			DATE:

FULL NAME OF INVENTOR	FAMILY NAME ROE	FIRST GIVEN NAME DAVID	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY ROCKWOOD	STATE OR COUNTRY CANADA	COUNTRY OF CITIZENSHIP CANADA
POST OFFICE ADDRESS	POST OFFICE ADDRESS 100 INKERMAN STREET, #7	CITY ROCKWOOD	STATE & ZIP CODE/COUNTRY ONTARIO CANADA N0B 2K0
INVENTOR'S SIGNATURE:			DATE:

FULL NAME OF INVENTOR	FAMILY NAME LU	FIRST GIVEN NAME LINDA	SECOND GIVEN NAME FUNG
RESIDENCE & CITIZENSHIP	CITY Scarborough	STATE OR COUNTRY CANADA	COUNTRY OF CITIZENSHIP CANADA
POST OFFICE ADDRESS	POST OFFICE ADDRESS 63 Fawndale Crescent	CITY Scarborough	STATE & ZIP CODE/COUNTRY ONTARIO CANADA M1W 2X3
INVENTOR'S SIGNATURE:			DATE:

FULL NAME OF INVENTOR	FAMILY NAME THORNBACK	FIRST GIVEN NAME JOHN	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY North Stewkley	STATE OR COUNTRY Great Britain	COUNTRY OF CITIZENSHIP Great Britain
POST OFFICE ADDRESS	POST OFFICE ADDRESS The Thatched Cottage 88 High Street	CITY North Stewkley	STATE & ZIP CODE/COUNTRY Bedfordshire, Great Britain LU7 OEW
INVENTOR'S SIGNATURE:			DATE:

FULL NAME OF INVENTOR	FAMILY NAME POLLOCK	FIRST GIVEN NAME CATHERINE	SECOND GIVEN NAME M.
RESIDENCE & CITIZENSHIP	CITY Toronto	STATE OR COUNTRY Ontario	COUNTRY OF CITIZENSHIP Canada
POST OFFICE ADDRESS	POST OFFICE ADDRESS 187 Berry Rd., #16 ^{cp} 11/06/01	CITY Toronto	STATE & ZIP CODE/COUNTRY Ontario, Canada, M8Y1W9
INVENTOR'S SIGNATURE:			DATE: 11/06/01

20310-1041500



PATENT DECLARATION
FILE NO. 7126-2
DE&Y File No.:2274-020133
PAGE 1 OF 3

DECLARATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or the below named inventors are the original, first and joint inventors (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled **IMMOBILIZED LABELLING COMPOUNDS AND METHODS**, the Specification of which:

- ☐ Is attached hereto.
☒ was filed on August 10, 2001 as Application Serial No. 09/913401.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims.

I acknowledge the duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability of the subject matter claimed in this application, as "materiality" is defined in Title 37, Code of Federal Regulations, § 1.56.

I hereby claim priority benefits under Title 35, United States Code, § 119 and/or § 365 of any foreign application(s) for patent or inventor's certificate, PCT international application(s), and United States provisional application(s), listed below and have also identified below any foreign application for patent or inventor's certificate, PCT international application, or United States provisional application, having a filing date before that of the application on which priority is claimed:

PRIORITY APPLICATION(S)

			Priority Claimed
60/120,561	UNITED STATES	17 February 1999	Yes
(Number)	(Country)	(Date Filed)	Yes/No
PCT/CA99/00700	CANADA	3 August 1999	Yes
(Number)	(Country)	(Date Filed)	Yes/No

I hereby direct that all correspondence and telephone calls be addressed to Barbara S. Kitchell, Akerman, Senterfitt & Eidson, P.A., Fourth Floor, 222 Lakeview Avenue, P.O. Box 3188, West Palm Beach, FL 33402-3188, (561) 653-5123.

I HEREBY DECLARE THAT ALL STATEMENTS MADE OF MY OWN KNOWLEDGE ARE TRUE AND THAT ALL STATEMENTS MADE ON INFORMATION AND BELIEF ARE BELIEVED TO BE TRUE; AND FURTHER THAT THESE STATEMENTS WERE MADE WITH THE KNOWLEDGE THAT WILLFUL FALSE STATEMENTS AND THE LIKE SO MADE ARE PUNISHABLE BY FINE OR IMPRISONMENT, OR BOTH UNDER SECTION 1001 OF TITLE 18 OF THE UNITED STATES CODE AND THAT SUCH WILLFUL FALSE STATEMENTS MAY JEOPARDIZE THE VALIDITY OF THE APPLICATION OR ANY PATENT ISSUED THEREON.



FULL NAME OF INVENTOR	FAMILY NAME POLLAK	FIRST GIVEN NAME ALFRED	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY	STATE OR COUNTRY SLOVENIA	COUNTRY OF CITIZENSHIP SLOVENIA
POST OFFICE ADDRESS	POST OFFICE ADDRESS MURNIKOVA 8A, 1000 LUJBANA	CITY	STATE & ZIP CODE/COUNTRY SLOVENIA
INVENTOR'S SIGNATURE:			DATE:


FULL NAME OF INVENTOR	FAMILY NAME ROE	FIRST GIVEN NAME DAVID	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY ROCKWOOD	STATE OR COUNTRY CANADA	COUNTRY OF CITIZENSHIP CANADA
POST OFFICE ADDRESS	POST OFFICE ADDRESS 100 INKERMANN STREET, #7	CITY ROCKWOOD	STATE & ZIP CODE/COUNTRY ONTARIO CANADA N0B 2K0
INVENTOR'S SIGNATURE:			DATE:

FULL NAME OF INVENTOR	FAMILY NAME LU	FIRST GIVEN NAME LINDA	SECOND GIVEN NAME FUNG
RESIDENCE & CITIZENSHIP	CITY Scarborough	STATE OR COUNTRY CANADA	COUNTRY OF CITIZENSHIP CANADA
POST OFFICE ADDRESS	POST OFFICE ADDRESS 63 Fawndale Crescent	CITY Scarborough	STATE & ZIP CODE/COUNTRY ONTARIO CANADA M1W 2X3
INVENTOR'S SIGNATURE:			DATE:

FULL NAME OF INVENTOR	FAMILY NAME THORNBACK	FIRST GIVEN NAME JOHN	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY London	STATE OR COUNTRY Great Britain	COUNTRY OF CITIZENSHIP Great Britain
POST OFFICE ADDRESS	POST OFFICE ADDRESS MRC Clinical Sciences Centre, Imperial College School of Medicine, Hammersmith Campus, Du Cane Road	CITY London	STATE & ZIP CODE/COUNTRY Great Britain W12 0NN
INVENTOR'S SIGNATURE:			DATE: 20 th Sept 2002

FULL NAME OF INVENTOR	FAMILY NAME POLLOCK	FIRST GIVEN NAME CATHERINE	SECOND GIVEN NAME M.
RESIDENCE & CITIZENSHIP	CITY Toronto	STATE OR COUNTRY Ontario	COUNTRY OF CITIZENSHIP Canada
POST OFFICE ADDRESS	POST OFFICE ADDRESS 59 Station Road Apt. C	CITY Toronto	STATE & ZIP CODE/COUNTRY Ontario Canada M8V 2R2
INVENTOR'S SIGNATURE:			DATE:

2024-10-10 10:10:10



DECLARATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or the below named inventors are the original, first and joint inventors (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled **IMMOBILIZED LABELLING COMPOUNDS AND METHODS**, the Specification of which:

- ☐ Is attached hereto.
☒ was filed on August 10, 2001 as Application Serial No. 09/913401.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims.

I acknowledge the duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability of the subject matter claimed in this application, as "materiality" is defined in Title 37, Code of Federal Regulations, § 1.56.

I hereby claim priority benefits under Title 35, United States Code, § 119 and/or § 365 of any foreign application(s) for patent or inventor's certificate, PCT international application(s), and United States provisional application(s), listed below and have also identified below any foreign application for patent or inventor's certificate, PCT international application, or United States provisional application, having a filing date before that of the application on which priority is claimed:

PRIORITY APPLICATION(S)

			Priority Claimed
			Yes
(Number)	(Country)	(Date Filed)	Yes/No
60/120,561	UNITED STATES	17 February 1999	Yes
PCT/CA99/00700	CANADA	3 August 1999	Yes
(Number)	(Country)	(Date Filed)	Yes/No

I hereby direct that all correspondence and telephone calls be addressed to Barbara S. Kitchell, Akerman, Senterfitt & Eidson, P.A., Fourth Floor, 222 Lakeview Avenue, P.O. Box 3188, West Palm Beach, FL 33402-3188, (561) 653-5123.

I HEREBY DECLARE THAT ALL STATEMENTS MADE OF MY OWN KNOWLEDGE ARE TRUE AND THAT ALL STATEMENTS MADE ON INFORMATION AND BELIEF ARE BELIEVED TO BE TRUE; AND FURTHER THAT THESE STATEMENTS WERE MADE WITH THE KNOWLEDGE THAT WILLFUL FALSE STATEMENTS AND THE LIKE SO MADE ARE PUNISHABLE BY FINE OR IMPRISONMENT, OR BOTH UNDER SECTION 1001 OF TITLE 18 OF THE UNITED STATES CODE AND THAT SUCH WILLFUL FALSE STATEMENTS MAY JEOPARDIZE THE VALIDITY OF THE APPLICATION OR ANY PATENT ISSUED THEREON.



FULL NAME OF INVENTOR	FAMILY NAME POLLAK	FIRST GIVEN NAME ALFRED	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY	STATE OR COUNTRY SLOVENIA	COUNTRY OF CITIZENSHIP SLOVENIA
POST OFFICE ADDRESS	POST OFFICE ADDRESS MURNIKOVA 8A, 1000 LUJBJANA	CITY	STATE & ZIP CODE/COUNTRY SLOVENIA
INVENTOR'S SIGNATURE:			DATE:

FULL NAME OF INVENTOR	FAMILY NAME ROE	FIRST GIVEN NAME DAVID	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY ROCKWOOD	STATE OR COUNTRY CANADA	COUNTRY OF CITIZENSHIP CANADA
POST OFFICE ADDRESS	POST OFFICE ADDRESS 100 INKERMAN STREET, #7	CITY ROCKWOOD	STATE & ZIP CODE/COUNTRY ONTARIO CANADA N0B 2KO
INVENTOR'S SIGNATURE:			DATE:

FULL NAME OF INVENTOR	FAMILY NAME LU	FIRST GIVEN NAME LINDA	SECOND GIVEN NAME FUNG LING
RESIDENCE & CITIZENSHIP	CITY Scarborough	STATE OR COUNTRY CANADA	COUNTRY OF CITIZENSHIP CANADA
POST OFFICE ADDRESS	POST OFFICE ADDRESS 63 Fawndale Crescent	CITY Scarborough	STATE & ZIP CODE/COUNTRY ONTARIO CANADA M1W 2X3
INVENTOR'S SIGNATURE:			DATE: 4 Oct 2001

FULL NAME OF INVENTOR	FAMILY NAME THORNBACK	FIRST GIVEN NAME JOHN	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY North Stewkley	STATE OR COUNTRY Great Britain	COUNTRY OF CITIZENSHIP Great Britain
POST OFFICE ADDRESS	POST OFFICE ADDRESS The Thatched Cottage 88 High Street	CITY North Stewkley	STATE & ZIP CODE/COUNTRY Bedfordshire, Great Britain LU7 OEW
INVENTOR'S SIGNATURE:			DATE:

FULL NAME OF INVENTOR	FAMILY NAME POLLOCK	FIRST GIVEN NAME CATHERINE	SECOND GIVEN NAME M.
RESIDENCE & CITIZENSHIP	CITY Toronto	STATE OR COUNTRY Ontario	COUNTRY OF CITIZENSHIP Canada
POST OFFICE ADDRESS	POST OFFICE ADDRESS 59 Station Road Apt. C	CITY Toronto	STATE & ZIP CODE/COUNTRY Ontario Canada M8V 2R2
INVENTOR'S SIGNATURE:			DATE:

2024-10-10 14:50



PATENT DECLARATION
FILE NO. 7126-2
DE&Y File No.: 2274-020133
PAGE 1 OF 3

DECLARATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or the below named inventors are the original, first and joint inventors (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled **IMMOBILIZED LABELLING COMPOUNDS AND METHODS**, the Specification of which:

- ☐ Is attached hereto.
☒ was filed on August 10, 2001 as Application Serial No. 09/913401.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims.

I acknowledge the duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability of the subject matter claimed in this application, as "materiality" is defined in Title 37, Code of Federal Regulations, § 1.56

I hereby claim priority benefits under Title 35, United States Code, § 119 and/or § 365 of any foreign application(s) for patent or inventor's certificate, PCT international application(s), and United States provisional application(s), listed below and have also identified below any foreign application for patent or inventor's certificate, PCT international application, or United States provisional application, having a filing date before that of the application on which priority is claimed.

PRIORITY APPLICATION(S)

			Priority Claimed
60/120,561	UNITED STATES	17 February 1999	Yes
(Number)	(Country)	(Date Filed)	Yes/No
PCT/CA99/00700	CANADA	3 August 1999	Yes
(Number)	(Country)	(Date Filed)	Yes/No

I hereby direct that all correspondence and telephone calls be addressed to Barbara S. Kitchell, Akerman, Senterfitt & Eidson, P.A., Fourth Floor, 222 Lakeview Avenue, P.O. Box 3188, West Palm Beach, FL 33402-3188, (561) 653-5123.

I HEREBY DECLARE THAT ALL STATEMENTS MADE OF MY OWN KNOWLEDGE ARE TRUE AND THAT ALL STATEMENTS MADE ON INFORMATION AND BELIEF ARE BELIEVED TO BE TRUE; AND FURTHER THAT THESE STATEMENTS WERE MADE WITH THE KNOWLEDGE THAT WILLFUL FALSE STATEMENTS AND THE LIKE SO MADE ARE PUNISHABLE BY FINE OR IMPRISONMENT, OR BOTH UNDER SECTION 1001 OF TITLE 18 OF THE UNITED STATES CODE AND THAT SUCH WILLFUL FALSE STATEMENTS MAY JEOPARDIZE THE VALIDITY OF THE APPLICATION OR ANY PATENT ISSUED THEREON



FULL NAME OF INVENTOR	FAMILY NAME POLLAK	FIRST GIVEN NAME ALFRED	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY	STATE OR COUNTRY SLOVENIA	COUNTRY OF CITIZENSHIP SLOVENIA
POST OFFICE ADDRESS	POST OFFICE ADDRESS MURNIKOVA 8A, 1000 LUJBANA	CITY	STATE & ZIP CODE/COUNTRY SLOVENIA
INVENTOR'S SIGNATURE:			DATE:

FULL NAME OF INVENTOR	FAMILY NAME ROE	FIRST GIVEN NAME DAVID	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY ONTARIO	STATE OR COUNTRY CANADA	COUNTRY OF CITIZENSHIP BRITISH CANADA <i>DeRoe.</i>
POST OFFICE ADDRESS	POST OFFICE ADDRESS 143 Academy Place, RR #2	CITY Toronto	STATE & ZIP CODE/COUNTRY Ontario, Canada, NOB 2K0
INVENTOR'S SIGNATURE:	<i>X DeRoe.</i>		DATE: <i>X Nov. 14/2001</i>

FULL NAME OF INVENTOR	FAMILY NAME LU	FIRST GIVEN NAME LINDA	SECOND GIVEN NAME FUNG
RESIDENCE & CITIZENSHIP	CITY Scarborough	STATE OR COUNTRY CANADA	COUNTRY OF CITIZENSHIP CANADA
POST OFFICE ADDRESS	POST OFFICE ADDRESS 63 Fawndale Crescent	CITY Scarborough	STATE & ZIP CODE/COUNTRY ONTARIO CANADA M1W 2X3
INVENTOR'S SIGNATURE:			DATE:

FULL NAME OF INVENTOR	FAMILY NAME THORNBACK	FIRST GIVEN NAME JOHN	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY North Stewkley	STATE OR COUNTRY Great Britain	COUNTRY OF CITIZENSHIP Great Britain
POST OFFICE ADDRESS	POST OFFICE ADDRESS The Thatched Cottage 88 High Street	CITY North Stewkley	STATE & ZIP CODE/COUNTRY Bedfordshire, Great Britain LU7 OEW
INVENTOR'S SIGNATURE:			DATE:

FULL NAME OF INVENTOR	FAMILY NAME POLLOCK	FIRST GIVEN NAME CATHERINE	SECOND GIVEN NAME M.
RESIDENCE & CITIZENSHIP	CITY Toronto	STATE OR COUNTRY Ontario	COUNTRY OF CITIZENSHIP Canada
POST OFFICE ADDRESS	POST OFFICE ADDRESS 187 Berry Rd.	CITY Toronto	STATE & ZIP CODE/COUNTRY Ontario, Canada, M8Y1W9
INVENTOR'S SIGNATURE:			DATE:

2025-10-10 10:46:50



PATENT DECLARATION
FILE NO. 7126-2
DE&Y File No.:2274-020133
PAGE 1 OF 3

DECLARATION

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or the below named inventors are the original, first and joint inventors (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled **IMMOBILIZED LABELLING COMPOUNDS AND METHODS**, the Specification of which:

- ☐ Is attached hereto.
☒ was filed on August 10, 2001 as Application Serial No. 09/913401.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims.

I acknowledge the duty to disclose to the Patent and Trademark Office all information known to me to be material to patentability of the subject matter claimed in this application, as "materiality" is defined in Title 37, Code of Federal Regulations, § 1.56.

I hereby claim priority benefits under Title 35, United States Code, § 119 and/or § 365 of any foreign application(s) for patent or inventor's certificate, PCT international application(s), and United States provisional application(s), listed below and have also identified below any foreign application for patent or inventor's certificate, PCT international application, or United States provisional application, having a filing date before that of the application on which priority is claimed:

PRIORITY APPLICATION(S)

			Priority Claimed
60/120,561	UNITED STATES	17 February 1999	Yes
(Number)	(Country)	(Date Filed)	Yes/No
PCT/CA99/00700	CANADA	3 August 1999	Yes
(Number)	(Country)	(Date Filed)	Yes/No

I hereby direct that all correspondence and telephone calls be addressed to Barbara S. Kitchell, Akerman, Senterfitt & Eidson, P.A., Fourth Floor, 222 Lakeview Avenue, P.O. Box 3188, West Palm Beach, FL 33402-3188, (561) 653-5123.

I HEREBY DECLARE THAT ALL STATEMENTS MADE OF MY OWN KNOWLEDGE ARE TRUE AND THAT ALL STATEMENTS MADE ON INFORMATION AND BELIEF ARE BELIEVED TO BE TRUE; AND FURTHER THAT THESE STATEMENTS WERE MADE WITH THE KNOWLEDGE THAT WILLFUL FALSE STATEMENTS AND THE LIKE SO MADE ARE PUNISHABLE BY FINE OR IMPRISONMENT, OR BOTH UNDER SECTION 1001 OF TITLE 18 OF THE UNITED STATES CODE AND THAT SUCH WILLFUL FALSE STATEMENTS MAY JEOPARDIZE THE VALIDITY OF THE APPLICATION OR ANY PATENT ISSUED THEREON.



100

FULL NAME OF INVENTOR	FAMILY NAME POLLAK	FIRST GIVEN NAME ALFRED	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY	STATE OR COUNTRY SLOVENIA SKX	COUNTRY OF CITIZENSHIP SLOVENIA
POST OFFICE ADDRESS	POST OFFICE ADDRESS MURNIKOVA 8A, 1000 LUJBIANA	CITY	STATE & ZIP CODE/COUNTRY SLOVENIA
INVENTOR'S SIGNATURE:			DATE: Sep 18, 2001

200

FULL NAME OF INVENTOR	FAMILY NAME ROE	FIRST GIVEN NAME DAVID	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY	STATE OR COUNTRY CANADA CAX	COUNTRY OF CITIZENSHIP CANADA
POST OFFICE ADDRESS	POST OFFICE ADDRESS 100 INKERMANN STREET, #7	CITY ROCKWOOD	STATE & ZIP CODE/COUNTRY ONTARIO CANADA N0B 2K0
INVENTOR'S SIGNATURE:			DATE:

300

FULL NAME OF INVENTOR	FAMILY NAME LU	FIRST GIVEN NAME LINDA	SECOND GIVEN NAME FUNG
RESIDENCE & CITIZENSHIP	CITY Scarborough	STATE OR COUNTRY CANADA CAX	COUNTRY OF CITIZENSHIP CANADA
POST OFFICE ADDRESS	POST OFFICE ADDRESS 63 Fawndale Crescent	CITY Scarborough	STATE & ZIP CODE/COUNTRY ONTARIO CANADA M1W 2X3
INVENTOR'S SIGNATURE:			DATE:

400

FULL NAME OF INVENTOR	FAMILY NAME THORNBACK	FIRST GIVEN NAME JOHN	SECOND GIVEN NAME
RESIDENCE & CITIZENSHIP	CITY North Stewkley	STATE OR COUNTRY Great Britain SBN	COUNTRY OF CITIZENSHIP Great Britain
POST OFFICE ADDRESS	POST OFFICE ADDRESS The Thatched Cottage 88 High Street	CITY North Stewkley	STATE & ZIP CODE/COUNTRY Bedfordshire, Great Britain LU7 OEW
INVENTOR'S SIGNATURE:			DATE:

FULL NAME OF INVENTOR	FAMILY NAME <u>POLLOCK</u>	FIRST GIVEN NAME <u>CATHERINE</u>	SECOND GIVEN NAME <u>M.</u>
RESIDENCE & CITIZENSHIP	CITY <u>Toronto</u>	STATE OR COUNTRY <u>Ontario</u> <i>CAX</i>	COUNTRY OF CITIZENSHIP <u>Canada</u>
POST OFFICE ADDRESS	POST OFFICE ADDRESS <u>59 Station Road Apt. C</u>	CITY <u>Toronto</u>	STATE & ZIP CODE/COUNTRY <u>Ontario Canada M8V 2R2</u>
INVENTOR'S SIGNATURE:			DATE:

50

20970-1047660



09/91 3401
Rec'd PCT/PTO 16 JAN 2002

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:

Alfred Pollack
David Rose
Linda Fung Lu
John Thornback
Catherine M. Pollock

Group Art Unit:

Serial No.: 09/913,401

Examiner: Not Assigned

Date Filed: August 10, 2001 1

Atty. Dkt. No.: 7126-2

For: IMMOBILIZED LABELLING COMPOUNDS AND METHODS

**ELECTION UNDER 37 C.F.R. §§ 3.71 AND 3.73
AND POWER OF ATTORNEY**

Commissioner of Patents
Washington, D.C. 20231

Sir:

The undersigned, being Assignee of record of the entire interest in the above-identified application by virtue of an assignment recorded in the United States Patent and Trademark Office as set forth below, hereby elects, under 37 C.F.R. § 3.71, to prosecute the application to the exclusion of the inventors.

The Assignee hereby revokes any previous Powers of Attorney and appoints:

J. Rodman Steele, Jr.	Reg. No. <u>25,931</u>
Gregory A. Nelson	Reg. No. <u>30,577</u>
Joseph W. Bain	Reg. No. <u>34,290</u>
Robert J. Sacco	Reg. No. <u>35,667</u>
Mark D. Passler	Reg. No. <u>40,764</u>
Stanley A. Kim	Reg. No. <u>42,730</u>
Steven M. Greenberg	Reg. No. <u>44,725</u>
Neil R. Jetter	Reg. No. <u>46,803</u>
Larry G. Brown	Reg. No. <u>45,834</u>
Kevin T. Cuenot	Reg. No. <u>46,283</u>
Pablo Meles	Reg. No. <u>33,739</u>
Raynarldo Whitty	Reg. No. <u>47,176</u>
Barbara S. Kitchell	Reg. No. <u>33,928</u>

14

each an attorney or agent of the firm Akerman, Senterfitt & Eidson, P.A., as its attorney or agent for so long as they remain with such firm, with full power of substitution and revocation to prosecute the application, to make alterations and amendments therein, to transact all business in the Patent and Trademark Office in connection therewith, and to receive any Letters Patent, and for one year after issuance of such Letters Patent to file any request for a certificate of correction that may be deemed appropriate.

Pursuant to 37 C.F.R. § 3.73, the undersigned has reviewed the evidentiary documents, specifically the Assignment to BRACCO INTERNATIONAL B.V., referenced below, and certifies that to the best of my knowledge and belief, title remains in the name of the Assignee.

Please direct all communications as follows:

BARBARA S. KITCHELL
AKERMAN, SENTERFITT & EIDSON, P.A.
Fourth Floor
222 Lakeview Avenue
P.O. Box 3188
West Palm Beach, Florida 33402-3188
(561) 653-5123

ASSIGNEE:
BRACCO INTERNATIONAL B.V.

By:

Name: Gea van Estrik

Tom Herpschleb

Title: Proxy Holder

Proxy Holder

Date: 1 October 2001

ASSIGNMENT:

- ☐ Concurrently filed
- ☐ Previously recorded
- Date: